arXiv:1204.2690v1 [math.RT] 12 Apr 2012

Tensor products of unipotent characters of general linear groups over finite fields

Emmanuel Letellier
Université de Caen
`letellier.emmanuel@math.unicaen.fr`

March 2, 2013

Abstract

Given unipotent characters $\mathcal{U}_1, \dots, \mathcal{U}_k$ of $\mathrm{GL}_n(\mathbb{F}_q)$, we prove that $\langle \mathcal{U}_1 \otimes \cdots \otimes \mathcal{U}_k, 1\rangle$ is a polynomial in q with non-negative integer coefficients (this was observed for $n \leq 8$ and $k = 3$ by Hiss-Lübeck-Mattig [8]). We study the degree of this polynomial and give a necessary and sufficient condition in terms of the representation theory of symmetric groups and root systems for this polynomial to be non-zero.

Contents

1 The main results **1**
1.1 Generic case . . . 2
1.1.1 Connection with quiver varieties . . . 3
1.1.2 Connection with character varieties . . . 3
1.2 Unipotent case . . . 4

2 Preliminaries **6**
2.1 Log and Exp . . . 6
2.2 Partitions, types . . . 7
2.3 Littlewood-Richardson coefficients . . . 8
2.4 A technical result . . . 10
2.5 Cauchy function . . . 12
2.6 Harcos inequality . . . 13

3 Tensor products of unipotent characters **14**
3.1 Irreducible characters of unipotent type . . . 14
3.2 Comet-shaped quivers . . . 14
3.3 The generic case . . . 16
3.4 The unipotent case . . . 18
3.5 Example . . . 23

1 The main results

Recall that the complex unipotent characters of $\mathrm{GL}_n(\mathbb{F}_q)$ are naturally parameterized by the irreducible characters of the symmetric group $\mathfrak{S}_n$ and therefore by the partitions of n. For a partition μ of n we put $\mathcal{U}_\mu$

the corresponding unipotent character. Under our parametrization, the trivial character of GL_n is $\mathcal{U}_{(n^1)}$ and the Steinberg character is $\mathcal{U}_{(1^n)}$.

Fix an integer $g \geq 0$ and consider $\mathcal{E} : \mathrm{GL}_n(\mathbb{F}_q) \to \mathbb{C}$, $x \mapsto q^{g \dim C_{\mathrm{GL}_n(\overline{\mathbb{F}}_q)}(x)}$. If $g = 1$, this is the character of the representation of $\mathrm{GL}_n(\mathbb{F}_q)$ in the group algebra $\mathbb{C}[\mathfrak{gl}_n(\mathbb{F}_q)]$ where GL_n acts on $\mathfrak{gl}_n$ by conjugation.

Using the inner product formula

$$\langle f, h \rangle = \langle f, h \rangle_{\mathrm{GL}_n(\mathbb{F}_q)} = \frac{1}{|\mathrm{GL}_n(\mathbb{F}_q)|} \sum_{x \in \mathrm{GL}_n} f(x)\overline{h(x)}$$

which holds for any two class functions $f, h : \mathrm{GL}_n(\mathbb{F}_q) \to \mathbb{C}$, it is not difficult to see, using the character table of $\mathrm{GL}_n(\mathbb{F}_q)$ due to Green [4], that for any multi-partition $\boldsymbol{\mu} = (\mu^1, \ldots, \mu^k)$ of n, there exists a polynomial $U_{\boldsymbol{\mu}}(t) \in \mathbb{Q}[t]$ such that for any finite field $\mathbb{F}_q$, we have

$$U_{\boldsymbol{\mu}}(q) = \left\langle \mathcal{E} \otimes \mathcal{U}_{\mu^1} \otimes \cdots \otimes \mathcal{U}_{\mu^k}, 1 \right\rangle.$$

The aim of this paper is to study the polynomials $U_{\boldsymbol{\mu}}(t)$. They were computed for $k = 3$, $g = 0$ and $1 \leq n \leq 8$ by F. Lübeck [12]. Some results of this paper can be easily observed in these tables.

1.1 Generic case

In order to state our main theorem on the polynomials $U_{\boldsymbol{\mu}}(q)$ we need to introduce an other class of polynomials $V_{\boldsymbol{\mu}}(q)$. We say that a tuple $(\mathcal{X}_1, \ldots, \mathcal{X}_k)$ of irreducible characters of $\mathrm{GL}_n(\mathbb{F}_q)$ is of *type* $\boldsymbol{\mu} = (\mu^1, \ldots, \mu^k)$ if for all $i = 1, \ldots, k$, there exists a linear character $\alpha_i : \mathbb{F}_q^\times \to \mathbb{C}^\times$ such that

$$\mathcal{X}_i = (\alpha_i \circ \det) \cdot \mathcal{U}_{\mu^i}.$$

Such a tuple is said to be *generic* if the linear character $\alpha_1 \alpha_2 \cdots \alpha_k$ is of order n.

In [11, §6.10.6] (for a review see also §3.3) we define polynomials $V_{\boldsymbol{\mu}}(t) \in \mathbb{Q}[t]$ for any multi-partition $\boldsymbol{\mu}$ and prove that for any finite field $\mathbb{F}_q$ and any generic tuple $(\mathcal{X}_1, \ldots, \mathcal{X}_k)$ of irreducible characters of $\mathrm{GL}_n(\mathbb{F}_q)$ of type $\boldsymbol{\mu}$ we have

$$V_{\boldsymbol{\mu}}(q) = \langle \mathcal{E} \otimes \mathcal{X}_1 \otimes \cdots \otimes \mathcal{X}_k, 1 \rangle.$$

From a multi-partition $\boldsymbol{\mu} = (\mu^1, \ldots, \mu^k)$, we define a comet-shaped graph $\Gamma_{\boldsymbol{\mu}}$ together with a dimension vector $\mathbf{v}_{\boldsymbol{\mu}}$ as in §3.2. We then denote by $\Phi(\Gamma_{\boldsymbol{\mu}})$ the associated root system as defined in [9].

Put

$$d_{\boldsymbol{\mu}} := n^2(2g - 2 + k) - \sum_{i,j} (\mu^i_j)^2 + 2 = 2 - {}^t\mathbf{v}_{\boldsymbol{\mu}} \mathbf{C}_{\boldsymbol{\mu}} \mathbf{v}_{\boldsymbol{\mu}}$$

where $\mathbf{C}_{\boldsymbol{\mu}}$ is the Cartan matrix of $\Gamma_{\boldsymbol{\mu}}$ and $\mu^i = (\mu^i_1, \ldots, \mu^i_{r_i})$ with $\mu^i_1 \geq \mu^i_2 \geq \cdots \geq \mu^i_{r_i}$.

Theorem 1.1.1. *(i) The polynomial* $V_{\boldsymbol{\mu}}(t)$ *is non-zero if and only if* $\mathbf{v}_{\boldsymbol{\mu}} \in \Phi(\Gamma_{\boldsymbol{\mu}})$. *Moroever* $V_{\boldsymbol{\mu}}(t) = 1$ *if and only if* $\mathbf{v}_{\boldsymbol{\mu}}$ *is a real root.*
(ii) If non-zero, $V_{\boldsymbol{\mu}}(t)$ *is a monic polynomial of degree* $d_{\boldsymbol{\mu}}/2$ *with non-negative integer coefficients.*

Remark 1.1.2. In [11] we defined the notion of generic tuples $(\mathcal{X}_1, \ldots, \mathcal{X}_k)$ for any types (not necessarily unipotent) of irreducible characters $\mathcal{X}_1, \ldots, \mathcal{X}_k$ of $\mathrm{GL}_n(\mathbb{F}_q)$. Among these generic tuples, we defined a subclass whose elements are called *admissible generic tuples*. We then proved that if $(\mathcal{X}_1, \ldots, \mathcal{X}_k)$ is admissible then the inner product $\langle \mathcal{E} \otimes \mathcal{X}_1 \otimes \cdots \otimes \mathcal{X}_k, 1 \rangle$ can be expressed as the Poincaré polynomial (for intersection cohomology) of a certain quiver variety, from which we prove a statement analogous to Theorem 1.1.1. Unfortunately, generic tuples of irreducible characters of unipotent type are never admissible and so we can not use the results of [11] to prove Theorem 1.1.1.

1.1.1 Connection with quiver varieties

Consider a generic tuple $(C_1, \ldots, C_k)$ of regular semisimple adjoint orbits of $\mathfrak{gl}_n(\mathbb{C})$ (see proof of Theorem 3.3.2 for the definition of generic tuples) and consider the space $\mathcal{V}$ of tuples of matrices

$$(A_1, \ldots, A_g, B_1, \ldots, B_g, X_1, \ldots, X_k) \in \mathfrak{gl}_n(\mathbb{C})^{2g} \times C_1 \times \cdots \times C_k$$

which satisfy the equation

$$[A_1, B_1] + \cdots + [A_g, B_g] + X_1 + \cdots + X_k = 0.$$

Put

$$Q := \mathcal{V} /\!/ \mathrm{GL}_n = \mathrm{Spec}\left(\mathbb{C}[\mathcal{V}]^{\mathrm{GL}_n}\right)$$

where GL_n acts diagonally by conjugation on $\mathcal{V}$. The variety Q is non-singular and the quotient map $\mathcal{V} \to Q$ is a principal PGL_n-bundle in the étale topology. Denote by $H_c^i(Q, \mathbb{C})$ the compactly supported cohomology of Q. Recall (see for instance [5]) that $H_c^i(Q, \mathbb{C}) = 0$ when i is odd.

We can define an action ρ^i of k copies $\mathbb{S}_n := \mathfrak{S}_n \times \cdots \times \mathfrak{S}_n$ of the symmetric group $\mathfrak{S}_n$ on $H_c^{2i}(Q, \mathbb{C})$. This is a particular case of Weyl group actions on cohomology of quiver varieties constructed and studied by many authors including Nakajima [17] [18], Lusztig [13], Maffei [15]. The construction of the Weyl group action given in [11] does not apply here (we can only construct the action of some relative Weyl groups which are finite subgroups of $\mathbb{S}_n$).

For a partition λ, denote by χ^λ the irreducible character of the symmetric group $\mathfrak{S}_n$ associated with λ as in [14]. Following the strategy of [7] (see proof of Theorem 3.3.2 for more details) we can show that for any multipartition $\boldsymbol{\mu} = (\mu^1, \ldots, \mu^k)$ of n we have

$$V_{\boldsymbol{\mu}}(t) = q^{-d/2} \sum_i \left\langle \chi^{\boldsymbol{\mu}'}, \rho^i \right\rangle_{\mathbb{S}_n} t^i \tag{1.1.1}$$

where d is the dimension of Q, $\boldsymbol{\mu}'$ denotes the dual multi-partition of $\boldsymbol{\mu}$ and $\chi^{\boldsymbol{\mu}}$ is the irreducible character $\chi^{\mu^1} \otimes \cdots \otimes \chi^{\mu^k}$ of $\mathbb{S}_n$. Formula (1.1.1) implies the positivity of the coefficients of $V_{\boldsymbol{\mu}}(t)$. Theorem 1.1.1(i) together with Formula (1.1.1) provides a nice criterion in terms of roots for the appearance or not of an irreducible character of $\mathbb{S}_n$ in $\rho^* := \bigoplus_i \rho^i$.

1.1.2 Connection with character varieties

Let us recall the conjectural interpretation of the polynomials $V_{\boldsymbol{\mu}}(t)$ in terms of Poincaré polynomial of character varieties [11, §1.3].

For a partition λ of n let us denote by C_λ the unipotent conjugacy class of $\mathrm{GL}_n(\mathbb{C})$ whose size of Jordan blocks is given by the dual partition λ' of λ.

For a multi-partition $\boldsymbol{\mu} = (\mu^1, \ldots, \mu^k)$ of n, put

$$\overline{C}_{\boldsymbol{\mu}} := \mathrm{GL}_n^{2g} \times \overline{C}_{\mu^1} \times \cdots \times \overline{C}_{\mu^k}.$$

Fix primitive n-th root of unity ζ and consider the space $\mathcal{Z}_{\boldsymbol{\mu}}$ of tuples

$$(A_1, \ldots, A_g, B_1, \ldots, B_g, X_1, \ldots, X_k) \in \overline{C}_{\boldsymbol{\mu}}$$

which satisfy the equation

$$\prod_{i=1}^{g} (A_i, B_i) \prod_{j=1}^{k} X_i = \zeta \cdot I_n$$

where I_n is the identity matrix and (A, B) is the commutator $ABA^{-1}B^{-1}$. Put

$$\mathcal{M}_{\mu} := \mathcal{Z}_{\mu}//\mathrm{GL}_n = \mathrm{Spec}\left(\mathbb{C}[\mathcal{Z}_{\mu}]^{\mathrm{GL}_n}\right)$$

where GL_n acts diagonally by conjugation on $\mathcal{Z}_{\mu}$. By Saito [19] the compactly supported intersection cohomology $IH^i_c(\mathcal{M}_{\mu}, \mathbb{C})$ is endowed with a mixed Hodge structure. Denote by $\{ih_c^{r,s;k}(\mathcal{M}_{\mu})\}_{r,s,k}$ the corresponding mixed Hodge numbers and consider the pure part

$$PP_c(\mathcal{M}_{\mu}, t) := \sum_s ih_c^{s,s;2s}(\mathcal{M}_{\mu})t^s$$

of the mixed Poincaré polynomial. Then we have the following conjecture [11, Conjecture 1.3.2]

Conjecture 1.1.3.

$$V_{\mu}(t) = t^{-d_{\mu}/2}PP_c(\mathcal{M}_{\mu}, t).$$

1.2 Unipotent case

In order to see the relation between the two polynomials $U_{\mu}(t)$ and $V_{\mu}(t)$, we need to introduce some notations. Consider k separate sets $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_k$ of infinitely many variables and denote by $\Lambda(\mathbf{x}_1, \ldots, \mathbf{x}_k) = \Lambda(\mathbf{x}_1) \otimes_{\mathbb{Z}} \cdots \otimes_{\mathbb{Z}} \Lambda(\mathbf{x}_k)$ the ring of functions separately symmetric in each set $\mathbf{x}_1, \ldots, \mathbf{x}_k$, and put $\Lambda = \mathbb{Q}(t) \otimes_{\mathbb{Z}} \Lambda(\mathbf{x}_1, \ldots, \mathbf{x}_k)$. For a multi-partition $\boldsymbol{\mu} = (\mu^1, \ldots, \mu^k)$, we define $s_{\boldsymbol{\mu}} \in \Lambda$ by

$$s_{\boldsymbol{\mu}} := s_{\mu^1}(\mathbf{x}_1) \cdots s_{\mu^k}(\mathbf{x}_k)$$

where for a partition λ we denote by $s_\lambda(\mathbf{x}_i) \in \Lambda(\mathbf{x}_i)$ the corresponding Schur symmetric function as in [14].

Denote by $\mathcal{P}$ the set of all partitions including the unique partition 0 of 0 and denote by $\overline{\mathcal{P}}$ the set of multi-partitions $\boldsymbol{\mu} = (\mu^1, \ldots, \mu^k) \in \mathcal{P}^k$ with $|\mu^1| = |\mu^2| = \cdots = |\mu^k| =: |\boldsymbol{\mu}|$. We denote by $\mathcal{P}_n$ and $\overline{\mathcal{P}}_n$ the subsets of partitions of size n.

We prove the following result (see Proposition 3.4.1).

Proposition 1.2.1. *We have*

$$\mathrm{Exp}\left(\sum_{\boldsymbol{\mu}\in\overline{\mathcal{P}}-\{0\}} V_{\boldsymbol{\mu}}(t)s_{\boldsymbol{\mu}}T^{|\boldsymbol{\mu}|}\right) = 1 + \sum_{\boldsymbol{\mu}\in\overline{\mathcal{P}}-\{0\}} U_{\boldsymbol{\mu}}(t)s_{\boldsymbol{\mu}}T^{|\boldsymbol{\mu}|}$$

where $\mathrm{Exp} : T\Lambda[[T]] \to 1 + T\Lambda[[T]]$ *is the plethystic exponential.*

Our strategy to study the polynomials $U_{\mu}(t)$ is to use the above proposition together with the properties of $V_{\mu}(t)$.

Consider now a total ordering $\geq$ on the set of all partitions. Denote by $\mathbf{T}_n^o$ the set of non-increasing sequences of partitions $\alpha^1\alpha^2\cdots\alpha^r$ such that $\sum_{i=1}^r |\alpha^i| = n$. We will write the elements $\omega^o \in \mathbf{T}_n^o$ in the form $(\alpha^1)^{n_1}(\alpha^2)^{n_2}\cdots(\alpha^s)^{n_s}$ with $\alpha^1 > \alpha^2 > \cdots > \alpha^s$ and with n_i the multiplicity of α^i in ω^o. We then put

$$\mathfrak{S}_{\omega^o} := \prod_{i=1}^s (\mathfrak{S}_{|\alpha^i|})^{n_i}, \qquad H_{\omega^o} := \bigotimes_{i=1}^s T^{n_i}H_{\alpha^i}$$

where for a partition λ, H_λ is an irreducible $\mathbb{C}[\mathfrak{S}_{|\lambda|}]$-module with character χ^λ, and where $T^mV := V\otimes\cdots\otimes V$, with V repeated m times. For a partition μ of n and a type $\omega^o \in \mathbf{T}_n^o$ define

$$\mathbb{C}^{\mu}_{\omega^o} := \mathrm{Hom}_{\mathfrak{S}_n}\left(\mathrm{Ind}_{\mathfrak{S}_{\omega^o}}^{\mathfrak{S}_n}(H_{\omega^o}), H_\mu\right)$$

where for an inclusion of finite groups $H \subset K$, we denote by Ind_H^K is the usual induction functor $V \mapsto \mathbb{C}[K] \otimes_{\mathbb{C}[H]} V$ from the category of left $\mathbb{C}[H]$-modules into the category of left $\mathbb{C}[K]$-modules. In [11, §6] we constructed an action of the group $W_{\omega^o} := \mathfrak{S}_{n_1}\times\cdots\times\mathfrak{S}_{n_s}$ on the $\mathbb{C}$-vector space $\mathbb{C}^{\mu}_{\omega^o}$ and we proved that given a partition $\nu_i = (d_{i,1},\dots,d_{i,r_i})$ of n_i for all $i = 1,\dots,s$, the coordinates of

$$s_{\alpha^1}(\mathbf{x}^{d_{1,1}})\cdots s_{\alpha^1}(\mathbf{x}^{d_{1,r_1}})s_{\alpha^2}(\mathbf{x}^{d_{2,1}})\cdots s_{\alpha^2}(\mathbf{x}^{d_{2,r_2}})\cdots s_{\alpha^s}(\mathbf{x}^{d_{s,1}})\cdots s_{\alpha^s}(\mathbf{x}^{d_{s,r_s}})$$

in the basis of Schur symmetric functions $\{s_\mu\}_\mu$ equal Trace $\left(w\,|\,\mathbb{C}^{\mu}_{\omega^o}\right)$ where $w = (w_1,\dots,w_s) \in W_{\omega^o}$ with w_i in the conjugacy class of $\mathfrak{S}_{n_i}$ corresponding to the partition ν_i. Choose once for all a total ordering on the set of multi-partitions $\overline{\mathcal{P}}$ and denote by $\overline{\mathbf{T}}^o_n$ the set of non-increasing sequences $\boldsymbol{\alpha}_1^{n_1}\boldsymbol{\alpha}_2^{n_2}\cdots\boldsymbol{\alpha}_s^{n_s}$ such that $\boldsymbol{\alpha}_1 > \boldsymbol{\alpha}_2 > \cdots > \boldsymbol{\alpha}_s$ and

$$\sum_{i=1}^{s} n_i|\boldsymbol{\alpha}_i| = n.$$

It will be also convenient in this paper to think of the element $\boldsymbol{\omega}^o = \boldsymbol{\alpha}_1^{n_1}\boldsymbol{\alpha}_2^{n_2}\cdots\boldsymbol{\alpha}_s^{n_s} \in \overline{\mathbf{T}}^o_n$ as a function $\boldsymbol{\omega}^o : \overline{\mathcal{P}} \to \mathbb{Z}_{\geq 0}$ with $\boldsymbol{\omega}^o(\boldsymbol{\alpha}_i) = n_i$ and $\boldsymbol{\omega}^o(\boldsymbol{\mu}) = 0$ if $\boldsymbol{\mu} \notin \{\boldsymbol{\alpha}_1,\dots,\boldsymbol{\alpha}_s\}$.

The total orderings on $\mathcal{P}$ and $\overline{\mathcal{P}}$ defines a natural map $\overline{\mathbf{T}}^o_n \to (\mathbf{T}^o_n)^k$, $\boldsymbol{\omega}^o \mapsto (\omega^o_1,\dots,\omega^o_k)$. For $\boldsymbol{\mu} = (\mu^1,\dots,\mu^k) \in \overline{\mathcal{P}}_n$ and $\boldsymbol{\omega}^o \in \mathbf{T}^o_n$, define

$$\mathbb{C}^{\boldsymbol{\mu}}_{\omega^o} := \bigotimes_{i=1}^{k} \mathbb{C}^{\mu^i}_{\omega^o_i}.$$

If $\boldsymbol{\omega}^o = \boldsymbol{\alpha}_1^{n_1}\boldsymbol{\alpha}_2^{n_2}\cdots\boldsymbol{\alpha}_s^{n_s}$, the group $W_{\omega^o} := \prod_{i=1}^s \mathfrak{S}_{n_i}$ acts on $\mathbb{C}^{\boldsymbol{\mu}}_{\omega^o_i}$ via its diagonal embedding in $W_{\omega^o_1}\times\cdots\times W_{\omega^o_k}$. We then define

$$\mathcal{R}_{\omega^o,\boldsymbol{\mu}} := \left\{(\tau^1,\dots,\tau^s) \in \mathcal{P}_{n_1}\times\cdots\times\mathcal{P}_{n_s} \;\middle|\; \left\langle H_{\tau^1}\otimes\cdots\otimes H_{\tau^s}, \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}\right\rangle_{W_{\omega^o}} \neq 0\right\}.$$

For a partition λ, we denote by $\ell(\lambda)$ its length. We can now state the main result of this paper.

Theorem 1.2.2. *Let $\boldsymbol{\mu} \in \overline{\mathcal{P}}_n$ with $n \geq 1$.*
(i) The polynomial $U_{\boldsymbol{\mu}}(t)$ has non-negative integer coefficients.
(ii) The polynomial $U_{\boldsymbol{\mu}}(t)$ is non-zero if and only if there exists $\boldsymbol{\omega}^o = \boldsymbol{\alpha}_1^{n_1}\boldsymbol{\alpha}_2^{n_2}\cdots\boldsymbol{\alpha}_s^{n_s} \in \overline{\mathbf{T}}^o_n$ and $(\tau^1,\dots,\tau^s) \in \mathcal{R}_{\omega^o,\boldsymbol{\mu}}$ such that

$$\ell(\tau^i) \leq V_{\alpha_i}(1) \tag{1.2.1}$$

for all $i = 1,\dots,s$.

By Theorem 1.1.1, the inequality (1.2.1) does not hold unless $\mathbf{v}_{\alpha_i}$ is a root of Γ_{α_i}. Denote by $\overline{\mathbf{T}}^o_{n+}$ the subset of $\overline{\mathbf{T}}^o_n$ of sequences $\boldsymbol{\alpha}_1^{n_1}\boldsymbol{\alpha}_2^{n_2}\cdots\boldsymbol{\alpha}_s^{n_s}$ with $\mathbf{v}_{\alpha_i} \in \Phi(\Gamma_{\alpha_i})$.

Corollary 1.2.3. *Let $\boldsymbol{\mu} \in \overline{\mathcal{P}}_n$ with $n \geq 1$. If there exists $\omega^o \in \overline{\mathbf{T}}^o_{n+}$ such that $\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1\right\rangle \neq 0$, then $U_{\boldsymbol{\mu}}(t) \neq 0$.*

Let $\boldsymbol{\omega}^o = \boldsymbol{\alpha}_1^{n_1}\boldsymbol{\alpha}_2^{n_2}\cdots\boldsymbol{\alpha}_s^{n_s} \in \overline{\mathbf{T}}^o_n$ and $\boldsymbol{\mu} = (\mu^1,\dots,\mu^k) \in \overline{\mathcal{P}}_n$.

Notice that if $\boldsymbol{\alpha}_i = ((1),\dots,(1))$ for all $i = 1,\dots,s$, then $W_{\omega^o} = \mathfrak{S}_n$ and $\mathbb{C}^{\boldsymbol{\mu}}_{\omega^o} = H_{\boldsymbol{\mu}}$.

Note also that if $n_1 = n_2 = \cdots = n_s$, then $W_{\omega^o} = 1$ and so

$$\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1\right\rangle = \dim \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}$$

is a product of Littlewood-Richardson coefficients. If moreover $s = 1$ and $n_1 = 1$, then $\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1\right\rangle = \delta_{\omega^o,\boldsymbol{\mu}}$.

In particular it follows from Corollary 1.2.3 that if $\mathbf{v}_\mu \in \Phi(\Gamma_\mu)$ or if $\langle H_{\boldsymbol{\mu}}, 1\rangle \neq 0$, then $U_{\boldsymbol{\mu}}(t) \neq 0$.

We can actually prove the following result (see Proposition 3.4.4 and Remark 3.4.5).

Proposition 1.2.4. *The term $\langle H_{\boldsymbol{\mu}}, 1\rangle$ contributes to the constant term of $U_{\boldsymbol{\mu}}(t)$.*

For $\boldsymbol{\mu} = (\mu^1, \dots, \mu^k)$ put

$$\delta(\boldsymbol{\mu}) = (2g-2+k)n - \sum_{i=1}^{k} \mu_1^i.$$

Then $\mathbf{v}_{\mu}$ is in the fundamental set of imaginary roots of Γ_{μ} if and only if $\delta(\boldsymbol{\mu}) \geq 0$. Note also that if $g \geq 1$, then $\delta(\boldsymbol{\mu}) \geq 0$ and so in this case $U_{\mu}(t)$ is always non-zero.

We prove the following theorem concerning the degree of $U_{\mu}(t)$.

Theorem 1.2.5. *(i) If* $\mathbf{v}_{\mu} \in \Phi(\Gamma_{\mu})$*, then the degree of* $U_{\mu}(t)$ *is at least* $d_{\mu}/2$*.*
(ii) If $\delta(\boldsymbol{\mu}) \geq 2$*, then the degree of* $U_{\mu}(t)$ *is exactly* $d_{\mu}/2$*.*
(iii) If $\delta(\boldsymbol{\mu}) \geq 3$ *or* $g = 0, k = 3$ *and* $\delta(\boldsymbol{\mu}) = 2$*, then* $U_{\mu}(t)$ *is monic.*

This theorem can be used to reduce the proof of results of the following kind to a finite number of checks (see proof of Corollary 3.4.10).

Corollary 1.2.6. *Let* St_n *denotes the Steinberg character of* $\mathrm{GL}_n(\mathbb{F}_q)$*. Then for all* $n \geq 1$*, the inner product* $\langle \mathrm{St}_n \otimes \mathrm{St}_n \otimes \mathrm{St}_n, 1\rangle$ *is a monic polynomial in* q *of degree* $\frac{1}{2}(n-1)(n-2)$*.*

2 Preliminaries

2.1 Log and Exp

Fix an integer $k > 0$. Consider k separate sets $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ of infinitely many variables and denote by $\Lambda(\mathbf{x}_1, \dots, \mathbf{x}_k) := \Lambda(\mathbf{x}_1) \otimes_{\mathbb{Z}} \cdots \otimes_{\mathbb{Z}} \Lambda(\mathbf{x}_k)$ the ring of functions separately symmetric in each set $\mathbf{x}_1, \dots, \mathbf{x}_k$. Put $\Lambda := \mathbb{Q}(t) \otimes_{\mathbb{Z}} \Lambda(\mathbf{x}_1, \dots, \mathbf{x}_k)$.

Consider

$$\psi_n : \Lambda[[T]] \to \Lambda[[T]],\ f(\mathbf{x}_1, \dots, \mathbf{x}_k; t, T) \mapsto f(\mathbf{x}_1^n, \dots, \mathbf{x}_k^n; t^n, T^n)$$

where we denote by $\mathbf{x}^d$ the set of variables $\{x_1^d, x_2^d, \dots\}$. The ψ_n are called the *Adams operations*.

Define $\Psi : T\Lambda[[T]] \to T\Lambda[[T]]$ by

$$\Psi(f) = \sum_{n \geq 1} \frac{\psi_n(f)}{n}.$$

Its inverse is given by

$$\Psi^{-1}(f) = \sum_{n \geq 1} \mu(n) \frac{\psi_n(f)}{n}$$

where μ is the ordinary Möbius function.

Following Getzler [3] we define $\mathrm{Log} : 1 + T\Lambda[[T]] \to T\Lambda[[T]]$ and its inverse $\mathrm{Exp} : T\Lambda[[T]] \to 1 + T\Lambda[[T]]$ as

$$\mathrm{Log}(f) = \Psi^{-1}(\log(f))$$

and

$$\mathrm{Exp}(f) = \exp(\Psi(f)).$$

Lemma 2.1.1. *Let* $f \in T\Lambda[[T]]$*. If* f *has coefficients in* $\mathbb{Z}[t] \otimes_{\mathbb{Z}} \Lambda(\mathbf{x}_1, \dots, \mathbf{x}_k) \subset \Lambda$*, then* $\mathrm{Exp}(f)$ *has also coefficients in* $\mathbb{Z}[t] \otimes_{\mathbb{Z}} \Lambda(\mathbf{x}_1, \dots, \mathbf{x}_k)$*.*

Proof. We could have defined Exp using the σ-operations instead of the ψ-operations in which case the above lemma becomes clear, see for instance [16] for more details. □

For $g \in \Lambda$ and $n \geq 1$ we put

$$g_n := \frac{1}{n} \sum_{d|n} \mu(d) \psi_{\frac{n}{d}}(g).$$

This is the Möbius inversion formula of $\psi_n(g) = \sum_{d|n} d \cdot g_d$.

We have the following lemma [16].

Lemma 2.1.2. *Let* $g \in \Lambda$ *and* $f_1, f_2 \in 1 + T\Lambda[[T]]$ *such that*

$$\log(f_1) = \sum_{d=1}^{\infty} g_d \cdot \log(\psi_d(f_2)).$$

Then

$$\mathrm{Log}(f_1) = g \cdot \mathrm{Log}(f_2).$$

2.2 Partitions, types

Denote by $\mathcal{P}$ the set of all partitions including the unique partition 0 of 0 and by $\mathcal{P}_n$ the subset of partitions of n. Partitions λ are denoted by $(\lambda_1, \lambda_2, \ldots, \lambda_r)$ with $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_r \geq 0$. We will sometimes write a partition λ as $(1^{m_1}, 2^{m_2}, \ldots, i^{m_i})$ where m_i denotes the multiplicity of i in λ. The *size* of λ is defined as $|\lambda| := \sum_i \lambda_i$. If $d > 0$ is an integer and $\lambda = (\lambda_1, \ldots, \lambda_r)$ a partition of n, then $d \cdot \lambda := (d\lambda_1, \ldots, d\lambda_r) \in \mathcal{P}$ is a partition of dn. We also define the sum $\lambda + \mu$ of two partitions $\lambda = (\lambda_1, \ldots, \lambda_r)$ and $\mu = (\mu_1, \ldots, \mu_s)$ as the partition $(\lambda_1 + \mu_1, \lambda_2 + \mu_2, \ldots)$. We consider on $\mathcal{P}_n$ the partial ordering $\trianglelefteq$ defined as follows. We have $\lambda \trianglelefteq \mu$ if for all i, $\lambda_1 + \cdots + \lambda_i \leq \mu_1 + \cdots + \mu_i$. We denote by $\overline{\mathcal{P}}$ the set of multi-partitions $\boldsymbol{\mu} = (\mu^1, \mu^2, \ldots, \mu^k) \in \mathcal{P}^k$ such that $|\mu^1| = |\mu^2| = \cdots = |\mu^k|$ and we extend in the obvious way the definitions of $d \cdot \boldsymbol{\mu}$ (with $d \in \mathbb{Z}_{\geq 0}$) and $\boldsymbol{\lambda} + \boldsymbol{\mu}$. We denote by $\overline{\mathcal{P}}_n$ the subset of multi-partitions in $\overline{\mathcal{P}}$ of size n. Finally we say that $\boldsymbol{\lambda} \trianglelefteq \boldsymbol{\mu}$ if and only if $\lambda^i \trianglelefteq \mu^i$ for all $i = 1, \ldots, k$.

We call *multi-type* a function $\boldsymbol{\omega} : \mathbb{Z}_{\geq 0} \times \overline{\mathcal{P}} \to \mathbb{Z}_{\geq 0}$ such that its support $S_{\boldsymbol{\omega}} := \{(d, \boldsymbol{\mu}) \,|\, \boldsymbol{\omega}(d, \boldsymbol{\mu}) \neq 0\}$ is finite and does not contain pairs of the form $(0, \boldsymbol{\mu})$ or $(d, 0)$. We denote by 0 the multi-type corresponding to the zero function. The *degrees* of a multi-type $\boldsymbol{\omega}$ are the integers d such that $(d, \boldsymbol{\mu}) \in S_{\boldsymbol{\omega}}$ for some $\boldsymbol{\mu} \in \overline{\mathcal{P}}$. If the degres of $\boldsymbol{\omega}$ are all equal to 1, we say that $\boldsymbol{\omega}$ is *split*. We call $|\boldsymbol{\omega}| := \sum_{(d,\boldsymbol{\mu}) \in S_{\boldsymbol{\omega}}} d \cdot |\boldsymbol{\mu}| \cdot \boldsymbol{\omega}(d, \boldsymbol{\mu})$ the *size* of $\boldsymbol{\omega}$. We denote by $\overline{\mathbf{T}}$ the set of all multi-types as above and by $\overline{\mathbf{T}}_n$ the subset of multi-types of size n. We use the notation $\mathbf{T}$ (resp. $\mathbf{T}_n$) instead of $\overline{\mathbf{T}}$ (resp. $\overline{\mathbf{T}}_n$) for $k = 1$, and call simply an element of $\mathbf{T}$ a *type*.

Note that we have a natural map $\overline{\mathbf{T}} \to \mathbf{T}^k$ as follows. If $\boldsymbol{\omega} \in \overline{\mathbf{T}}$, then for each $i = 1, \ldots, k$, we define its i-th coordinate $\omega_i : \mathbb{Z}_{\geq 0} \times \mathcal{P} \to \mathbb{Z}_{\geq 0}$ as $\omega_i(d, \mu) = \sum_{\boldsymbol{\mu}} \boldsymbol{\omega}(d, \boldsymbol{\mu})$ where the sum is over the elements $\boldsymbol{\mu} \in \overline{\mathcal{P}}$ whose i-th coordinate is μ.

Finally for $\boldsymbol{\omega} \in \overline{\mathbf{T}}$ we define the multi-partition $\boldsymbol{\omega}_+ \in \overline{\mathcal{P}}$ as

$$\boldsymbol{\omega}_+ := \sum_{(d, \boldsymbol{\mu}) \in S_{\boldsymbol{\omega}}} (d\, \boldsymbol{\omega}(d, \boldsymbol{\mu})) \cdot \boldsymbol{\mu}.$$

Given a family $\{a_{\boldsymbol{\mu}}\}_{\boldsymbol{\mu} \in \overline{\mathcal{P}}}$ of elements of Λ, we extend its definition to multi-types $\boldsymbol{\omega} \in \overline{\mathbf{T}}$ as

$$a_{\boldsymbol{\omega}} := \prod_{(d, \boldsymbol{\mu}) \in S_{\boldsymbol{\omega}}} \psi_d(a_{\boldsymbol{\mu}})^{\boldsymbol{\omega}(d, \boldsymbol{\mu})}.$$

For a multi-type $\boldsymbol{\omega} \in \overline{\mathbf{T}} - \{0\}$, define

$$C^o_{\boldsymbol{\omega}} := \begin{cases} \frac{\mu(d)}{d} (-1)^{r_{\boldsymbol{\omega}} - 1} \frac{(r_{\boldsymbol{\omega}} - 1)!}{\prod_{\boldsymbol{\mu}} \boldsymbol{\omega}(d, \boldsymbol{\mu})!} & \text{if there is no } (d', \boldsymbol{\mu}) \in S_{\boldsymbol{\omega}} \text{ with } d' \neq d. \\ 0 & \text{otherwise.} \end{cases}$$

where μ is the ordinary Möbius function and $r_\omega := \sum_{(d,\boldsymbol{\mu})\in S_\omega} \omega(d,\boldsymbol{\mu})$.

We have the following lemma [5, §2.3.3].

Lemma 2.2.1. *Let* $\{a_\mu\}_{\mu\in\overline{\mathcal{P}}}$ *be a family of elements of* Λ *with* $a_0 = 1$. *Then*

$$\operatorname{Log}\left(\sum_{\mu\in\overline{\mathcal{P}}} a_\mu T^{|\boldsymbol{\mu}|}\right) = \sum_{\omega\in\overline{\mathbf{T}}-\{0\}} C^o_\omega a_\omega T^{|\omega|}. \tag{2.2.1}$$

For a multi-type $\omega \in \overline{\mathbf{T}}$, define

$$A^o_\omega := \prod_{(r,\boldsymbol{\mu})\in S_\omega} \frac{1}{r^{\omega(r,\boldsymbol{\mu})}\omega(r,\boldsymbol{\mu})!}.$$

The following lemma is also straightforward.

Lemma 2.2.2. *Let* $\{a_\mu\}_{\mu\in\overline{\mathcal{P}}}$ *be a family of elements of* Λ *with* $a_0 = 1$. *Then*

$$\operatorname{Exp}\left(\sum_{\mu\in\overline{\mathcal{P}}-\{0\}} a_\mu T^{|\boldsymbol{\mu}|}\right) = \sum_{\omega\in\overline{\mathbf{T}}} A^o_\omega a_\omega T^{|\omega|}.$$

The formal power series $\sum_{n\geq 0} a_n T^n$ with $a_n \in \Lambda$ that we will consider in what follows will all have a_n homogeneous of degree n in the variables $\mathbf{x}_1, \ldots, \mathbf{x}_k$. Hence we will typically scale the variables of Λ by $1/T$ and eliminate T altogether.

2.3 Littlewood-Richardson coefficients

For a partition $\lambda \in \mathcal{P}$ we denote by $s_\lambda(\mathbf{x}) \in \Lambda(\mathbf{x})$ the corresponding Schur function. For a type $\omega \in \mathbf{T}$ and a partition $\mu \in \mathcal{P}$, define $c^\mu_\omega \in \mathbb{Z}$ by

$$s_\omega = \sum_{\mu \trianglelefteq \omega_+} c^\mu_\omega s_\mu.$$

Note that $c^\mu_\omega = 0$ unless $|\omega| = |\mu|$. If ω is split, then c^μ_ω is a so-called *Littlewood-Richardson coefficient*.

For an integer $n > 0$, we denote by $\mathfrak{S}_n$ the symmetric group in n letters.

For a finite dimensional $\mathbb{C}$-vector space V and an integer $d > 0$, we put $T^d V := V \otimes \cdots \otimes V$ with V repeated d times.

For a partition λ, we denote by H_λ an irreducible $\mathbb{C}[\mathfrak{S}_{|\lambda|}]$-module corresponding to the irreducible character χ^λ of $\mathfrak{S}_{|\lambda|}$. Here we use the same parametrization $\lambda \mapsto \chi^\lambda$ as in [14]; the trivial character of $\mathfrak{S}_n$ corresponds to the partition (n).

Define $\mathbf{T}^o$ as the set of functions $\boldsymbol{\omega}^o : \mathcal{P} \to \mathbb{Z}_{\geq 0}$ whose support $S_{\omega^o} := \{\mu \,|\, \omega^o(\mu) \neq 0\}$ is finite and does not contain the 0 element of $\mathcal{P}$.

Note that we have a natural map $\mathfrak{H} : \mathbf{T} \to \mathbf{T}^o$ that maps ω to the function ω^o defined by $\omega^o(\mu) = \sum_d d \cdot \omega(d,\mu)$.

Given a type $\omega^o \in \mathbf{T}^o$, we put

$$H_{\omega^o} := \bigotimes_{\mu\in S_{\omega^o}} T^{\omega^o(\mu)} H_\mu, \qquad \mathfrak{S}_{\omega^o} := \prod_{\mu\in S_{\omega^o}} (\mathfrak{S}_{|\mu|})^{\omega^o(\mu)}, \qquad W_{\omega^o} := \prod_{\mu\in S_{\omega^o}} \mathfrak{S}_{\omega^o(\mu)}.$$

The elements of the fiber $\mathfrak{H}^{-1}(\boldsymbol{\omega}^o)$ are in bijection with $\prod_{\mu\in S_{\omega^o}} \mathcal{P}_{\omega^o(\mu)}$ and so with the conjugacy classes of W_{ω^o}. For $\omega^o \in \mathbf{T}^o$ and $\mu \in \mathcal{P}$ we define

$$\mathbb{C}^\mu_{\omega^o} := \operatorname{Hom}_{\mathfrak{S}_{|\omega^o|}}\left(\operatorname{Ind}^{\mathfrak{S}_{|\omega^o|}}_{\mathfrak{S}_{\omega^o}}(H_{\omega^o}), H_\mu\right).$$

The normalizer $N_{\mathfrak{S}_{|\omega^o|}}(\mathfrak{S}_{\omega^o})$ of $\mathfrak{S}_{\omega^o}$ in $\mathfrak{S}_{|\omega^o|}$ acts on the set of representations of $\mathfrak{S}_{\omega^o}$ on the left as $\sigma \cdot \rho := \rho \circ \sigma^{-1}$. Then we have an isomorphism

$$W_{\omega^o} \simeq \left\{\sigma \in N_{\mathfrak{S}_{|\omega^o|}}(\mathfrak{S}_{\omega^o}) \,\middle|\, \sigma \cdot \rho_{\omega^o} \simeq \rho_{\omega^o}\right\} / \mathfrak{S}_{\omega^o}.$$

By [11, §6.2], the group W_{ω^o} acts on the space $\mathbb{C}^{\mu}_{\omega^o}$ and we have the following proposition.

Proposition 2.3.1. *For all $v \in W_{\omega^o}$ we have*

$$\mathrm{Tr}\left(v \,\middle|\, \mathbb{C}^{\mu}_{\omega^o}\right) = c^{\mu}_{\omega}$$

where $\omega \in \mathbf{T}$ is the element in the fiber $\mathfrak{H}^{-1}(\omega^o)$ which corresponds to the conjugacy class of v.

We now extend this proposition to the case of multi-partitions.

Given any family $\{a_\mu\}$ of symmetric functions indexed by partitions $\mu \in \mathcal{P}$ and a multi-partition $\boldsymbol{\mu} = (\mu^1, \ldots, \mu^k) \in \overline{\mathcal{P}}$ define

$$a_{\boldsymbol{\mu}} := a_{\mu^1}(\mathbf{x}_1) \cdots a_{\mu^k}(\mathbf{x}_k).$$

For a multi-type $\boldsymbol{\omega} \in \overline{\mathbf{T}}$ and a multi-partition $\boldsymbol{\mu} \in \overline{\mathcal{P}}$, we denote by $c^{\boldsymbol{\mu}}_{\boldsymbol{\omega}}$ the integer defined by

$$s_{\boldsymbol{\omega}} = \sum_{\boldsymbol{\mu} \trianglelefteq \boldsymbol{\omega}_+} c^{\boldsymbol{\mu}}_{\boldsymbol{\omega}} s_{\boldsymbol{\mu}}.$$

Remark 2.3.2. Denote by $(\omega_1, \ldots, \omega_k)$ the coordinates of $\boldsymbol{\omega}$ in $\mathbf{T}^k$. Then

$$s_{\boldsymbol{\omega}} = s_{\omega_1}(\mathbf{x}_1) \cdots s_{\omega_k}(\mathbf{x}_k),$$

and so we see that the coefficient $c^{\boldsymbol{\mu}}_{\boldsymbol{\omega}}$ is a product $c^{\mu^1}_{\omega_1} \cdots c^{\mu^k}_{\omega_k}$ of coefficients $c^{\mu^i}_{\omega_i}$ defined above where $\mu^1, \ldots, \mu^k$ are the coordinates of $\boldsymbol{\mu}$.

Define $\overline{\mathbf{T}}^o$ as the set of functions $\boldsymbol{\omega}^o : \overline{\mathcal{P}} \to \mathbb{Z}_{\geq 0}$ whose support $S_{\boldsymbol{\omega}^o} := \{\boldsymbol{\mu} \,|\, \boldsymbol{\omega}^o(\boldsymbol{\mu}) \neq 0\}$ is finite and does not contain the 0 element of $\overline{\mathcal{P}}$. If $k = 1$, then $\overline{\mathbf{T}}^o$ is simply $\mathbf{T}^o$ defined above. We then have a natural map $\overline{\mathbf{T}}^o \to (\mathbf{T}^o)^k$ that sends $\boldsymbol{\omega}^o$ to $(\omega^o_1, \ldots, \omega^o_k)$ with $\omega^o_i(\mu) := \sum_{\boldsymbol{\mu}} \boldsymbol{\omega}^o(\boldsymbol{\mu})$ where the sum is over the elements $\boldsymbol{\mu} \in \overline{\mathcal{P}}$ whose i-th coordinate is μ. We call ω^o_i the i-th coordinate of $\boldsymbol{\omega}^o$.

Given a multi-type $\boldsymbol{\omega}^o \in \overline{\mathbf{T}}^o$, we put

$$W_{\boldsymbol{\omega}^o} := \prod_{\boldsymbol{\mu} \in S_{\boldsymbol{\omega}^o}} \mathfrak{S}_{\boldsymbol{\omega}^o(\boldsymbol{\mu})}.$$

For all $i = 1, \ldots, k$, the group $W_{\boldsymbol{\omega}^o}$ is a subgroup of $W_{\omega^o_i}$.

Consider the map $\overline{\mathfrak{H}} : \overline{\mathbf{T}} \to \overline{\mathbf{T}}^o$ that maps $\boldsymbol{\omega}$ to the function $\boldsymbol{\omega}^o$ defined by $\boldsymbol{\omega}^o(\boldsymbol{\mu}) = \sum_d d \cdot \boldsymbol{\omega}(d, \boldsymbol{\mu})$. The elements of the fiber $\overline{\mathfrak{H}}^{-1}(\boldsymbol{\omega}^o)$ are then in bijection with $\prod_{\boldsymbol{\mu} \in S_{\boldsymbol{\omega}^o}} \mathcal{P}_{\boldsymbol{\omega}^o(\boldsymbol{\mu})}$ and so with the conjugacy classes of $W_{\boldsymbol{\omega}^o}$.

For $\boldsymbol{\omega}^o \in \overline{\mathbf{T}}^o$ with coordinates $(\omega^o_1, \ldots, \omega^o_k)$ and $\boldsymbol{\mu} = (\mu^1, \ldots, \mu^k) \in \overline{\mathcal{P}}$ we define

$$\mathbb{C}^{\boldsymbol{\mu}}_{\boldsymbol{\omega}^o} := \bigotimes_{i=1}^{k} \mathbb{C}^{\mu^i}_{\omega^o_i}$$

The group $W_{\omega^o_1} \times \cdots \times W_{\omega^o_k}$ acts on $\mathbb{C}^{\boldsymbol{\mu}}_{\boldsymbol{\omega}^o}$ and so does the group $W_{\boldsymbol{\omega}^o}$ via its diagonal embedding in $W_{\omega^o_1} \times \cdots \times W_{\omega^o_k}$.

The following proposition is a consequence of Proposition 2.3.1 and Remark 2.3.2.

Proposition 2.3.3. *For all $v \in W_{\boldsymbol{\omega}^o}$ we have*

$$\mathrm{Tr}\left(v \,\middle|\, \mathbb{C}^{\boldsymbol{\mu}}_{\boldsymbol{\omega}^o}\right) = c^{\boldsymbol{\mu}}_{\boldsymbol{\omega}}$$

where $\boldsymbol{\omega} \in \overline{\mathbf{T}}$ is the element in the fiber $\overline{\mathfrak{H}}^{-1}(\boldsymbol{\omega}^o)$ which corresponds to the conjugacy class of v.

2.4 A technical result

Assume given a family $\{V_\mu(t)\}_{\mu\in\overline{\mathcal{P}}}$ of polynomials in $\mathbb{Z}[t]$ indexed by $\overline{\mathcal{P}}$. Let $\{U_\mu(t)\}_{\mu\in\overline{\mathcal{P}}-\{0\}}$ be the family defined by

$$\mathrm{Exp}\left(\sum_{\mu\in\overline{\mathcal{P}}-\{0\}} V_\mu(t)s_\mu\right) = 1 + \sum_{\mu\in\overline{\mathcal{P}}-\{0\}} U_\mu(t)s_\mu.$$

By Lemma 2.1.1 we have $U_\mu(t)\in\mathbb{Z}[t]$.

The aim of this section is to study the properties of the polynomials $U_\mu(t)\in\mathbb{Z}[t]$ in terms of those of $V_\mu(t)$.

We have

$$\begin{aligned}
U_\mu(t) &= \left\langle \sum_{\omega\in\overline{\mathbf{T}}} A^o_\omega V_\omega(t)s_\omega, s_\mu \right\rangle\\
&= \sum_{\omega\in\overline{\mathbf{T}}} A^o_\omega V_\omega(t)c^\mu_\omega\\
&= \sum_{\omega^o\in\overline{\mathbf{T}}^o}\ \sum_{\omega\in\overline{\mathfrak{H}}^{-1}(\omega^o)} A^o_\omega V_\omega(t)c^\mu_\omega\\
&= \sum_{\omega^o\in\overline{\mathbf{T}}^o} W^{\omega^o}_\mu(t)
\end{aligned}$$

where

$$W^{\omega^o}_\mu(t) := \sum_{\omega\in\overline{\mathfrak{H}}^{-1}(\omega^o)} A^o_\omega V_\omega(t)c^\mu_\omega.$$

Now fix $\omega^o\in\overline{\mathbf{T}}^o$. Write $S_{\omega^o}=\{\alpha_1,\dots,\alpha_s\}$ and put $n_i:=\omega^o(\alpha_i)$.

Recall that the elements of $\overline{\mathfrak{H}}^{-1}(\omega^o)$ are naturally parameterized by the set $\mathcal{P}_{n_1}\times\cdots\times\mathcal{P}_{n_s}$. If $\omega\in\overline{\mathfrak{H}}^{-1}(\omega^o)$ corresponds to $(\lambda^1,\dots,\lambda^s)\in\mathcal{P}_{n_1}\times\cdots\times\mathcal{P}_{n_s}$, then $A^o_\omega = z_{\lambda^1}\cdots z_{\lambda^s}$ where for a partition $\lambda=(1^{m_1},2^{m_2},\dots)$ we put

$$z_\lambda := \prod_{i\geq 1} i^{m_i}\cdot m_i!.$$

Recall that z_λ is the cardinality of the centralizer in $\mathfrak{S}_{|\lambda|}$ of an element of type λ. For a partition λ, denote by $p_\lambda(\mathbf{x})\in\Lambda(\mathbf{x})$ the corresponding power symmetric function in the infinite set of variables $\mathbf{x}=\{x_1,x_2,\dots\}$. Then $p_1(\mathbf{x})=x_1+x_2+\cdots$. Let $\mathbf{y}_1,\dots,\mathbf{y}_s$ be s independent sets of infinitely many variables.

Consider

$$W^{\omega^o}_\mu(\mathbf{y}_1,\dots,\mathbf{y}_k) := \sum_{(\lambda^1,\dots,\lambda^s)\in\mathcal{P}_{n_1}\times\cdots\times\mathcal{P}_{n_s}} \frac{1}{z_{\lambda^1}\cdots z_{\lambda^s}}\, p_{\lambda^1}(\mathbf{y}_1)\cdots p_{\lambda^s}(\mathbf{y}_s)\,\mathrm{Tr}\left(v_{(\lambda^1,\dots,\lambda^s)}\,\middle|\,\mathbb{C}^\mu_{\omega^o}\right)$$

where $v_{(\lambda^1,\dots,\lambda^s)}\in W_{\omega^o}$ is a representative of the conjugacy class of W_{ω^o} corresponding to $(\lambda^1,\dots,\lambda^s)$.

Lemma 2.4.1. *Assume that for all $i=1,\dots,s$, the polynomial $V_{\alpha_i}(t)$ has non-negative integer coefficients, then for an appropriate specialization of the variables $\mathbf{y}_1,\dots,\mathbf{y}_s$, we have*

$$W^{\omega^o}_\mu(\mathbf{y}_1,\dots,\mathbf{y}_k) = W^{\omega^o}_\mu(t).$$

Proof. Since the coefficients of $V_{\alpha_i}(t)$ are non-negative, there is an appropriate specialization of $V_{\alpha_i}(1)$ variables in $\mathbf{y}_i = \{y_{i,1}, y_{i,2}, \dots\}$ into monomials t^i, with $i \geq 0$ (the other variables being specialized to 0) such that

$$p_1(\mathbf{y}_i) = V_{\alpha_i}(t).$$

If $\boldsymbol{\omega} \in \overline{\mathfrak{H}}^{-1}(\omega^o)$ correspond to $(\lambda^1, \dots, \lambda^s) \in \mathcal{P}_{n_1} \times \cdots \times \mathcal{P}_{n_s}$ where $\lambda^i = (\lambda^i_1, \lambda^i_2, \dots)$ then

$$\begin{aligned} V_{\omega}(t) &= \prod_{i=1}^{s} \prod_{j} V_{\alpha_i}\left(t^{\lambda^i_j}\right) \\ &= \prod_{i,j} p_1(\mathbf{y}_i^{\lambda^i_j}) \\ &= \prod_{i=1}^{s} p_{\lambda^i}(\mathbf{y}_i). \end{aligned}$$

□

Remark 2.4.2. By the discussion above Lemma 3.3.5, note that

$$\sum_{\omega \in \overline{\mathfrak{H}}^{-1}(\omega^o)} A^o_{\omega} c^{\boldsymbol{\mu}}_{\omega} = \left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle_{W_{\omega^o}}$$

and so if $V_{\alpha_i}(t) = 1$ for all $i = 1, \dots, s$, then $W^{\omega^o}_{\boldsymbol{\mu}}(q) = \left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle_{W_{\omega^o}}$.

We now decompose the character of the representation $W_{\omega^o} \to \mathrm{GL}(\mathbb{C}^{\boldsymbol{\mu}}_{\omega^o})$ as a sum of irreducible characters

$$\sum_{(\tau^1, \dots, \tau^s) \in \mathcal{P}_{n_1} \times \cdots \times \mathcal{P}_{n_s}} m_{(\tau^1, \dots, \tau^s)} \chi^{\tau^1} \cdots \chi^{\tau^s}.$$

We thus have

$$\begin{aligned} W^{\omega^o}_{\boldsymbol{\mu}}(\mathbf{y}_1, \dots, \mathbf{y}_k) &= \sum_{(\tau^1, \dots, \tau^s) \in \mathcal{P}_{n_1} \times \cdots \times \mathcal{P}_{n_s}} m_{(\tau^1, \dots, \tau^s)} \sum_{(\lambda^1, \dots, \lambda^s) \in \mathcal{P}_{n_1} \times \cdots \times \mathcal{P}_{n_s}} \frac{1}{z_{\lambda^1} \cdots z_{\lambda^s}} p_{\lambda^1}(\mathbf{y}_1) \cdots p_{\lambda^s}(\mathbf{y}_s) \chi^{\tau^1}_{\lambda^1} \cdots \chi^{\tau^s}_{\lambda^s}. \\ &= \sum_{(\tau^1, \dots, \tau^s) \in \mathcal{P}_{n_1} \times \cdots \times \mathcal{P}_{n_s}} m_{(\tau^1, \dots, \tau^s)} \prod_{i=1}^{s} \sum_{\lambda \in \mathcal{P}_{n_i}} \frac{1}{z_\lambda} p_\lambda(\mathbf{y}_i) \chi^{\tau^i}_{\lambda} \\ &= \sum_{(\tau^1, \dots, \tau^s) \in \mathcal{P}_{n_1} \times \cdots \times \mathcal{P}_{n_s}} m_{(\tau^1, \dots, \tau^s)} \, s_{\tau^1}(\mathbf{y}_1) \cdots s_{\tau^s}(\mathbf{y}_s). \end{aligned}$$

Now a Schur function s_λ decomposes as

$$s_\lambda = \sum_{\mu \trianglelefteq \lambda} K_{\lambda\mu} m_\mu$$

where m_μ is the monomial symmetric function associated with μ and $\{K_{\lambda\mu}\}_{\lambda,\mu}$ are the Kostka numbers which are non-negative integers. Hence

$$W^{\omega^o}_{\boldsymbol{\mu}}(\mathbf{y}_1, \dots, \mathbf{y}_k) = \sum_{(\tau^1, \dots, \tau^s) \in \mathcal{P}_{n_1} \times \cdots \times \mathcal{P}_{n_s}} f_{(\tau^1, \dots, \tau^s)} \, m_{\tau^1}(\mathbf{y}_1) \cdots m_{\tau^s}(\mathbf{y}_s), \tag{2.4.1}$$

for some $f_{(\tau^1, \dots, \tau^s)} \in \mathbb{Z}_{\geq 0}$.

Put

$$\mathcal{R}_{\omega^o,\mu} := \{(\tau^1,\dots,\tau^s)\in\mathcal{P}_{n_1}\times\cdots\times\mathcal{P}_{n_s}\,|\, m_{(\tau^1,\dots,\tau^s)}\neq 0\}.$$

For a partition λ, denote by $\ell(\lambda)$ its length.

Theorem 2.4.3. *Assume that for all $i=1,2,\dots,s$, the polynomial $V_{\alpha_i}(t)$ has non-negative integer coefficients. Then the polynomial $W_{\mu}^{\omega^o}(t)$ has non-negative integer coefficients. Moreover it is non-zero if and only if there exists a sequence $(\tau^1,\dots,\tau^s)\in\mathcal{R}_{\omega^o,\mu}$ such that for all $i=1,\dots,s$, we have*

$$\ell(\tau^i)\leq V_{\alpha_i}(1).$$

Proof. The assertion (i) follows from the fact that if we specialize the variables $\mathbf{y}_i$ according to Lemma 3.3.5 we see that the right hand side of Formula (2.4.1) is a polynomial in t with non-negative coefficients. We have

$$W_{\mu}^{\omega^o}(t)=\sum_{(\tau^1,\dots,\tau^s)\in\mathcal{P}_{n_1}\times\cdots\times\mathcal{P}_{n_s}} m_{(\tau^1,\dots,\tau^s)}\, s_{\tau^1}(\mathbf{y}_1)\cdots s_{\tau^s}(\mathbf{y}_s)\neq 0$$

if and only if there exists $(\tau_1,\dots,\tau_s)\in\mathcal{R}_{\omega^o,\mu}$ such that for all $i=1,\dots,s$ we have $s_{\tau^i}(\mathbf{y}_i)\neq 0$. But $s_{\tau^i}(\mathbf{y}_i)\neq 0$ if and only if there exists a partition $\lambda^i\trianglelefteq\tau^i$ such that $m_{\lambda^i}(\mathbf{y}_i)\neq 0$, i.e., such that $\ell(\lambda^i)\leq V_{\alpha_i}(1)$. Indeed, the integer $V_{\alpha_i}(1)$ is the number of variables in $\mathbf{y}_i$ that are specialized to a monomial q^i, the other variables being specialized to 0. We conclude by noticing that if $\lambda^i\trianglelefteq\tau^i$, then $\ell(\tau^i)\leq\ell(\lambda^i)$.

□

For simplicity choose a total ordering $\geq$ on $\overline{\mathcal{P}}$ and denote the elements of $\overline{\mathbf{T}}^o$ in the form $\omega^o=\alpha_1^{n_1}\alpha_2^{n_2}\cdots\alpha_s^{n_s}$ with $\alpha_1\geq\alpha_2\geq\cdots\geq\alpha_s$ and $\omega^o(\alpha_i)=n_i$.

Theorem 2.4.4. *Assume that the polynomials $V_\alpha(t)$, with $|\alpha|\leq n$, have non-negative integer coefficients. Then we have the following assertions.*
(i) For any $\mu\in\overline{\mathcal{P}}_n$, the polynomial $U_\mu(t)$ has non-negative integer coefficients.
(ii) The polynomial $U_\mu(t)$ is non-zero if and only if there exists $\omega^o=\alpha_1^{n_1}\alpha_2^{n_2}\cdots\alpha_s^{n_s}\in\overline{\mathbf{T}}^o$ and a sequence $(\tau^1,\dots,\tau^s)\in\mathcal{R}_{\omega^o,\mu}$ such that for all $i=1,\dots,s$, we have

$$\ell(\tau^i)\leq V_{\alpha_i}(1).$$

Proof. Follows from Theorem 2.4.3 and the fact that the polynomial $U_\mu(t)$ is non-zero if and only if there exists an $\omega^o\in\overline{\mathbf{T}}^o$ such that $W_\mu^{\omega^o}(t)\neq 0$. □

2.5 Cauchy function

Given a partition $\lambda\in\mathcal{P}_n$ and an integer $g\geq 0$, we define

$$\mathcal{H}_\lambda(t):=\frac{t^{g\langle\lambda,\lambda\rangle}}{a_\lambda(t)}. \tag{2.5.1}$$

where $a_\lambda(q)$ is the cardinality of the centralizer in $\mathrm{GL}_{|\lambda|}(\mathbb{F}_q)$ of a unipotent element with Jordan form of type λ.

For a partition λ, let $\tilde{H}_\lambda(\mathbf{x};t)=\sum_\nu\tilde{K}_{\nu\lambda}(t)s_\nu(\mathbf{x})\in\Lambda(\mathbf{x})\otimes_{\mathbb{Z}}\mathbb{Q}(t)$ where $\tilde{K}_{\nu\lambda}(t)$ are the modified Kostka polynomials [14, Chapter III, §7].

As in [5] we consider the function $\Omega(t)\in\Lambda[[T]]$ defined as

$$\Omega(t) = \Omega(\mathbf{x}_1, \ldots, \mathbf{x}_k; t) := \sum_{\lambda\in\mathcal{P}} \mathcal{H}_\lambda(t) \prod_{i=1}^{k} \tilde{H}_\lambda(\mathbf{x}_i; t).$$

When $g = 0$ and $k = 2$ this function was first considered by Garsia and Haiman [2]. Define $\mathbb{V}(t) \in T\Lambda[[T]]$ and $\mathbb{U}(t) \in 1 + T\Lambda[[T]]$ by

$$\mathbb{V}(t) := (t-1)\operatorname{Log}\Omega(t), \qquad \mathbb{U}(t) := \operatorname{Exp}\mathbb{V}(t)$$

2.6 Harcos inequality

We extend the definition of size of partitions to any sequence $\mathbf{x} = (x_1, \ldots, x_k)$ of non-negative integers as

$$|\mathbf{x}| := \sum_i x_i.$$

For two sequences $\mathbf{c} = (c_1, \ldots, c_r)$, $\mathbf{x} = (x_1, \ldots, x_s)$, with $c := \max_i c_i$ define

$$\sigma_{\mathbf{c}}(\mathbf{x}) := c|\mathbf{x}|^2 - |\mathbf{c}| \sum_i x_i^2.$$

Let us now state Harcos theorem [6, Appendix].

Theorem 2.6.1. *Let $r, s > 0$ be integers. For $i = 1, \ldots, s$, let $\mathbf{x}^i = (x^i_1, \ldots, x^i_r)$ be a sequence with non-negative numbers. Put $\mathbf{c} = (c_1, \ldots, c_r) := \sum_{i=1}^{s} \mathbf{x}^i$ and $c := \max_i c_i$. Then*

$$\sigma_{\mathbf{c}}(\mathbf{c}) \geq \sum_{i=1}^{s} \sigma_{\mathbf{c}}(\mathbf{x}^i).$$

Corollary 2.6.2. *Let $\alpha^1, \ldots, \alpha^s$ be s partitions and let μ be a partition of size $|\sum_i \alpha^i|$ such that $\mu \trianglelefteq \sum_i \alpha^i$. Then*

$$\sigma_\mu(\mu) \geq \sum_{i=1}^{s} \sigma_\mu(\alpha^i).$$

Proof. We have

$$\sigma_\mu(\mu) - \sum_{i=1}^{s} \sigma_\mu(\alpha^i) = \sigma_\mu(\mu) - \sum_{i=1}^{s} \left(\mu_1 |\alpha^i|^2 - |\mu| \sum_j (\alpha^i_j)^2 \right).$$

Since $\mu \trianglelefteq \sum_i \alpha^i$, we can find $\mathbf{x}^1, \ldots, \mathbf{x}^k$ such that

1) $\mu = \sum_i \mathbf{x}^i$,

2) for all $i = 1, \ldots, s$, $|\mathbf{x}^i| = |\alpha^i|$ and $\sum_j (x^i_j)^2 \leq \sum_j (\alpha^i_j)^2$.

Notices that the sequences $\mathbf{x}^i$ may not be partitions any more.

Hence

$$\sigma_\mu(\mu) - \sum_{i=1}^{s} \sigma_\mu(\alpha^i) \geq \sigma_\mu(\mu) - \sum_{i=1}^{s} \sigma_\mu(\mathbf{x}^i)$$

which is non-negative by Theorem 2.6.1.

$\square$

3 Tensor products of unipotent characters

On Λ we put $\langle\, ,\,\rangle := \prod_i \langle\, ,\,\rangle_i$ where $\langle\, ,\,\rangle_i$ denotes the Hall pairing on $\Lambda(\mathbf{x}_i)$ which makes the basis $\{s_\mu(\mathbf{x}_i)\}$ of Schur symmetric functions orthonormal.

3.1 Irreducible characters of unipotent type

To alleviate the notation, put $G := \mathrm{GL}_n(\mathbb{F}_q)$. Let $B \subset G$ be the upper triangular matrices and let $\mathbb{C}[G/B]$ be the $\mathbb{C}$-vector space with basis $G/B = \{gB\,|\,g \in G\}$. The group G acts on $\mathbb{C}[G/B]$ by left multiplication. Let us denote by $\mathrm{Ind}_B^G(1) : G \to \mathbb{C}, g \mapsto \mathrm{Trace}\,(g\,|\,\mathbb{C}[G/B])$ the character of the representation $G \to \mathrm{GL}\,(\mathbb{C}[G/B])$. The decomposition of $\mathrm{Ind}_B^G(1)$ as a sum of irreducible charaters of G reads

$$\mathrm{Ind}_B^G(1) = \sum_{\chi\in\mathrm{Irr}\,\mathfrak{S}_n} \chi(1)\cdot\mathcal{U}_\chi.$$

The irreducible characters $\{\mathcal{U}_\chi\}_\chi$ are called the *unipotent* characters of G. The character $\mathcal{U}_1$ is the trivial character of G and $\mathcal{U}_\epsilon$, where ϵ is the sign character of $\mathfrak{S}_n$, is the Steinberg character of G. For a partition λ of n, we put

$$\mathcal{U}_\lambda := \mathcal{U}_{\chi^\lambda}$$

so that the $\mathcal{U}_{(1^n)}$ is the Steinberg character and $\mathcal{U}_{(n^1)}$ is the trivial character.

We say that an irreducible character of G is of *unipotent type* if it is of the form $(\alpha\circ\det)\cdot\mathcal{U}_\lambda$ for some partition λ and some linear character $\alpha : \mathbb{F}_q^\times \to \mathbb{C}^\times$.

3.2 Comet-shaped quivers

Given a non-negative integer g and a k-tuple $\boldsymbol{\mu} = (\mu^1, \mu^2, \ldots, \mu^k) \in \overline{\mathcal{P}}_n$ with $n \geq 1$, we denote by $\Gamma_{\boldsymbol{\mu}}$ the *comet-shaped* quiver



with k legs of length $s_1, s_2, \ldots, s_k$ (where $s_i = \ell(\mu^i) - 1$) and with g loops at the central vertex. The multi-partition $\boldsymbol{\mu}$ defines also a dimension vector $\mathbf{v}_{\boldsymbol{\mu}}$ of $\Gamma_{\boldsymbol{\mu}}$ whose coordinates on the i-th leg are $(n, n-\mu^i_1, n-\mu^i_1-\mu^i_2, \ldots, n-\sum_{r=1}^{s_i}\mu^i_r)$.

Let $I_{\boldsymbol{\mu}} = \{0\}\cup\{[i,j]\,|\,1 \geq i \geq k,\ 1 \geq j \geq s_i\}$ be the set of vertices of $\Gamma_{\boldsymbol{\mu}}$ and let $\mathbf{C}_{\boldsymbol{\mu}} = (c_{ij})_{i,j}$ be the Cartan matrix of $\Gamma_{\boldsymbol{\mu}}$, namely

$$c_{ij} = \begin{cases} 2-2(\text{the number of edges joining } i \text{ to itself}) & \text{if } i = j \\ -(\text{the number of edges joining } i \text{ to } j) & \text{otherwise.} \end{cases}$$

Let $(\, ,\,)$ be the symmetric bilinear form on $\mathbb{Z}^{I_{\boldsymbol{\mu}}}$ defined by

$$(\mathbf{e}_i, \mathbf{e}_j) = c_{ij}.$$

where for $i \in I_{\boldsymbol{\mu}}$, we denote by $\mathbf{e}_i$ the root of $\Gamma_{\boldsymbol{\mu}}$ with all zero coordinates except for a 1 at the indicated vertex i. If there is no-edge loop at the vertex i, we say that $\mathbf{e}_i$ is a *fundamental root* [10, Chapter 1]. For a fundamental root $\mathbf{e}_i$ we define the associated fundamental reflection $s_i : \mathbb{Z}^{I_{\boldsymbol{\mu}}} \to \mathbb{Z}^{I_{\boldsymbol{\mu}}}$ by

$$s_i(\lambda) = \lambda - 2(\lambda, \mathbf{e}_i)\,\mathbf{e}_i$$

for all $\lambda \in \mathbb{Z}^I$. The group $W(\Gamma_{\boldsymbol{\mu}})$ generated by all fundamental reflections is called the Weyl group of $\Gamma_{\boldsymbol{\mu}}$. A vector $\mathbf{v} \in \mathbb{Z}^{I_{\boldsymbol{\mu}}}$ is called a *real root* of $\Gamma_{\boldsymbol{\mu}}$ if it is of the form $w(\mathbf{e}_i)$ for some fundamental root $\mathbf{e}_i$ and some $w \in W(\Gamma_{\boldsymbol{\mu}})$. Recall [10, Chapter 1] that the fundamental set $M(\Gamma_{\boldsymbol{\mu}})$ of imaginary roots is the set of vectors $\mathbf{v} \in (\mathbb{Z}_{\geq 0})^{I_{\boldsymbol{\mu}}} - \{0\}$ with connected support such that for all fundamental root $\mathbf{e}$, we have

$$(\mathbf{e}, \mathbf{v}) \leq 0.$$

The *imaginary roots* are the vectors $\mathbf{v} \in \mathbb{Z}^{I_{\boldsymbol{\mu}}}$ which are of the form $w(\delta)$ or $w(-\delta)$ for some $\delta \in M(\Gamma_{\boldsymbol{\mu}})$ and $w \in W(\Gamma_{\boldsymbol{\mu}})$.

Let $\Phi(\Gamma_{\boldsymbol{\mu}}) \subset \mathbb{Z}^I$ be the set of all roots (real and imaginary) of $\Gamma_{\boldsymbol{\mu}}$ and let $\Phi(\Gamma_{\boldsymbol{\mu}})^+ \subset (\mathbb{Z}_{\geq 0})^{I_{\boldsymbol{\mu}}}$ be the subset of positive roots.

For $\boldsymbol{\alpha} = (\alpha^1, \ldots, \alpha^k) \in \overline{\mathcal{P}}$, define

$$\delta(\boldsymbol{\alpha}) := (2g - 2 + k)n - \sum_{i=1}^{k} \alpha^i_1. \tag{3.2.1}$$

Proposition 3.2.1. *A dimension vector* $\mathbf{v}$ *of* $\Gamma_{\boldsymbol{\mu}}$ *is in* $M(\Gamma_{\boldsymbol{\mu}})$ *if and only if there exists* $\boldsymbol{\alpha} \in \overline{\mathcal{P}}$ *such that* $\mathbf{v} = \mathbf{v}_{\alpha}$ *and* $\delta(\boldsymbol{\alpha}) \geq 0$.

Proof. For all $i, j \geq 1$, we have

$$(\mathbf{v}, \mathbf{e}_{[i,j]}) = -\Big((v_{[i,j-1]} - v_{[i,j]}) - (v_{[i,j]} - v_{[i,j+1]})\Big)$$

where for convenience $[i, 0]$ denotes also the central vertex 0. We also have

$$\begin{aligned}(\mathbf{v}, \mathbf{e}_0) &= (2 - 2g)v_0 - \sum_{i=1}^{k} v_{[i,1]}\\ &= -\left((2g - 2 + k)v_0 - \sum_{i=1}^{k}(v_0 - v_{[i,1]})\right)\end{aligned}$$

For all $i = 1, \ldots, k$, put $\alpha^i_1 := v_0 - v_{[i,1]} \in \mathbb{Z}$ and $\alpha^i_j := v_{[i,j-1]} - v_{[i,j]}$ for all $j \geq 1$, and put $\alpha^i = (\alpha^i_1, \alpha^i_2, \ldots)$.

Then $\mathbf{v}$ is in the fundamental domain if and only if for all $i = 1, \ldots, k$, the tuple α^i is a partition and $\delta(\boldsymbol{\alpha}) \geq 0$ where $\boldsymbol{\alpha} = (\alpha^1, \ldots, \alpha^k)$. It is also clear that $\mathbf{v} = \mathbf{v}_{\alpha}$.

□

Note that if $g \geq 1$, then $\delta(\boldsymbol{\mu}) \geq 0$ and so $\mathbf{v}_{\boldsymbol{\mu}}$ is always an imaginary root.

Put

$$A_{\boldsymbol{\mu}}(t) := \left\langle \mathbb{V}(t), h_{\boldsymbol{\mu}} \right\rangle \tag{3.2.2}$$

where $\mathbb{V}(t)$ is as in §2.5 and $h_{\boldsymbol{\mu}} = h_{\mu^1}(\mathbf{x}_1) \cdots h_{\mu^k}(\mathbf{x}_k)$ denotes the complete symmetric function.

For $\boldsymbol{\mu} \in \overline{\mathcal{P}}$, put

$$d_{\boldsymbol{\mu}} := n^2(2g - 2 + k) - \sum_{i,j} (\mu^i_j)^2 + 2 = 2 - {}^t\mathbf{v}_{\boldsymbol{\mu}} \mathbf{C}_{\boldsymbol{\mu}} \mathbf{v}_{\boldsymbol{\mu}}. \tag{3.2.3}$$

Recall one of the main result of [6].

Theorem 3.2.2. *(i) For any finite field $\mathbb{F}_q$, the evaluation $A_\mu(q)$ counts the number of isomorphism classes of absolutely indecomposable representations of Γ_μ of dimension $\mathbf{v}_\mu$ over $\mathbb{F}_q$.*
(ii) If non-zero, $A_\mu(t)$ is a monic polynomial of degree $d_\mu/2$ with integer coefficients.
(iii) The polynomial $A_\mu(t)$ is non-zero if and only if $\mathbf{v}_\mu \in \Phi(\Gamma_\mu)$. Moreover $A_\mu(t) = 1$ if and only if $\mathbf{v}_\mu$ is a real root.

We recently proved in [7] that the coefficients of $A_\mu(t)$ are actually non-negative. The assertion (ii) follows from (i) using the results in [9, §1.15], and the assertion (iii) follows from (i) and [9, §1.10].

3.3 The generic case

Let $\boldsymbol{\mu} = (\mu^1, \dots, \mu^k) \in \overline{\mathcal{P}}_n$ with $n \geq 1$. A tuple $(\mathcal{X}_1, \dots, \mathcal{X}_k)$ of irreducible characters of G is said to be of type $\boldsymbol{\mu}$ if for each $i = 1, 2, \dots, k$, there exists a linear character $\alpha_i : \mathbb{F}_q^\times \to \mathbb{C}^\times$ such that

$$\mathcal{X}_i := (\alpha_i \circ \det) \cdot \mathcal{U}_{\mu^i}.$$

The tuple $(\mathcal{X}_1, \dots, \mathcal{X}_k)$ said to be *generic* if the size of the subgroup of $\mathrm{Irr}\,(\mathbb{F}_q^\times)$ generated by $\alpha_1 \cdots \alpha_k$ equals n (see [11, Definition 6.8.6]).

Fix an integer $g \geq 0$ and consider $\mathcal{E} : G \to \mathbb{C}$, $x \mapsto q^{g \dim C_{\mathrm{GL}_n}(x)}$. If $g = 1$, note that $\mathcal{E}$ is the character of the representation of G in the group algebra $\mathbb{C}[\mathfrak{g}]$ where G acts on $\mathfrak{g} := \mathfrak{gl}_n(\mathbb{F}_q)$ by conjugation.

Define

$$V_\mu(t) := \left\langle \mathbb{V}(t), s_\mu \right\rangle. \tag{3.3.1}$$

I.e., the $V_\mu(t)$ are defined by the identity

$$\sum_{\mu \in \overline{\mathcal{P}}} V_\mu(t) s_\mu = \mathbb{V}(t).$$

Recall the following theorem [11, §6.10.6].

Theorem 3.3.1. *For any generic tuple $(\mathcal{X}_1, \dots, \mathcal{X}_k)$ of type $\boldsymbol{\mu}$ we have*

$$\langle \mathcal{E} \otimes \mathcal{X}_1 \otimes \cdots \otimes \mathcal{X}_k, 1 \rangle_G = V_\mu(q) \tag{3.3.2}$$

By definition $V_\mu(t)$ is a rational function in t with rational coefficients and by the above theorem it is an integer for infinitely many values of t. Hence $V_\mu(t)$ is a polynomial in t with rational coefficients.

Theorem 3.3.2. *(1) The polynomial $V_\mu(t)$ is non-zero if and only if $\mathbf{v}_\mu \in \Phi(\Gamma_\mu)$. Moreover $V_\mu(t) = 1$ if and only if $\mathbf{v}_\mu$ is a real root.*
(2) If non-zero, the polynomial $V_\mu(t)$ is a monic polynomial of degree $d_\mu/2$ with non-negative integer coefficients.

For $\boldsymbol{\alpha} = (\alpha^1, \dots, \alpha^k), \boldsymbol{\beta} = (\beta^1, \dots, \beta^k) \in \overline{\mathcal{P}}_n$, say that $\boldsymbol{\alpha} \trianglelefteq \boldsymbol{\beta}$ if $\alpha^i \trianglelefteq \beta^i$ for all $i = 1, \dots, k$. We will need the following lemma.

Lemma 3.3.3. *If $\alpha, \boldsymbol{\beta} \in \overline{\mathcal{P}}_n$ are such that $\boldsymbol{\alpha} \trianglelefteq \boldsymbol{\beta}$ and $\boldsymbol{\alpha} \neq \boldsymbol{\beta}$ then $d_\beta < d_\alpha$.*

Proof. We need to see that for two partitions $\lambda = (\lambda_1, \dots, \lambda_r)$ and $\mu = (\mu_1, \dots, \mu_s)$ such that $\lambda \trianglelefteq \mu$ and $\lambda \neq \mu$, we have $\sum_{i=1}^r \lambda_i^2 < \sum_{i=1}^s \mu_i^2$. This follows from the formula

$$\sum_{i=1}^l \mu_i^2 - \sum_{i=1}^l \lambda_i^2 = (\mu_1-\lambda_1)(\mu_1-\mu_2+\lambda_1-\lambda_2) + (\mu_1+\mu_2-\lambda_1-\lambda_2)(\mu_2-\mu_3+\lambda_2-\lambda_3) + \cdots + \left(\sum_{i=1}^l \mu_i - \sum_{i=1}^l \lambda_i \right)(\mu_s+\lambda_s)$$

which is available for all l (with the convention that $\lambda_i = 0$ if $i > r$). □

Proof of Theorem 3.3.2. We have the following relations between Schur and complete symmetric functions [14, page 101]:

$$h_\lambda=\sum_{\mu\trianglerighteq\lambda}K'_{\lambda\mu}s_\mu,\qquad s_\mu=\sum_{\lambda\trianglerighteq\mu}K^*_{\mu\lambda}h_\lambda,$$

where $K=(K_{\lambda\mu})_{\lambda,\mu}$ is the matrix whose coefficients are Kostka numbers, $K'=(K'_{\lambda\mu})_{\lambda,\mu}$ is the transpose of K and $K^*=(K^*_{\lambda\mu})_{\lambda,\mu}$ is the transpose inverse of K. By Formulas (3.2.2) and (3.3.1) we have for any $\boldsymbol{\lambda},\boldsymbol{\mu}\in\overline{\mathcal{P}}-\{0\}$

$$A_{\boldsymbol{\lambda}}(t)=\sum_{\boldsymbol{\mu}\trianglerighteq\boldsymbol{\lambda}}K'_{\boldsymbol{\lambda}\boldsymbol{\mu}}V_{\boldsymbol{\mu}}(t),\qquad V_{\boldsymbol{\mu}}(t)=\sum_{\boldsymbol{\lambda}\trianglerighteq\boldsymbol{\mu}}K^*_{\boldsymbol{\mu}\boldsymbol{\lambda}}A_{\boldsymbol{\lambda}}(t),\tag{3.3.3}$$

where $K_{\boldsymbol{\lambda}\boldsymbol{\mu}}:=\prod_{i=1}^kK_{\lambda^i\mu^i}$. Assume that $\mathbf{v}_{\boldsymbol{\mu}}\in\Phi(\Gamma_{\boldsymbol{\mu}})$. By Theorem 3.2.2 the polynomial $A_{\boldsymbol{\mu}}(t)$ is monic of degree $d_{\boldsymbol{\mu}}/2$. Moreover $K^*_{\boldsymbol{\mu}\boldsymbol{\mu}}=1$ and by Lemma 3.3.3 the degree of the polynomials $A_{\boldsymbol{\lambda}}(t)$, with $\boldsymbol{\lambda}\trianglerighteq\boldsymbol{\mu}$, $\boldsymbol{\lambda}\neq\boldsymbol{\mu}$, are of degree strictly smaller than $d_{\boldsymbol{\mu}}/2$. Hence we deduce from the second formula (3.3.3) that $V_{\boldsymbol{\mu}}(t)$ is non-zero and is a monic polynomial of degree $d_{\boldsymbol{\mu}}/2$. Note that if $\mathbf{v}_{\boldsymbol{\mu}}$ is real, then $A_{\boldsymbol{\mu}}(t)=1$ and $A_{\boldsymbol{\lambda}}(t)=0$ if $\boldsymbol{\lambda}\trianglerighteq\boldsymbol{\mu},\boldsymbol{\lambda}\neq\boldsymbol{\mu}$ as $d_{\boldsymbol{\lambda}}<d_{\boldsymbol{\mu}}=0$, and so $V_{\boldsymbol{\mu}}(t)=1$. Assume now that $V_{\boldsymbol{\mu}}(t)\neq0$. Recall that $K_{\boldsymbol{\lambda}\boldsymbol{\mu}}$ are non-negative integers and that $K_{\boldsymbol{\mu}\boldsymbol{\mu}}=1$. Moreover, for all $\boldsymbol{\alpha}\in\overline{\mathcal{P}}-\{0\}$, the evaluation $V_{\boldsymbol{\alpha}}(q)$ of $V_{\boldsymbol{\alpha}}(t)$ at q is a non-negative integer by Theorem 3.3.1. Hence, by the first formula (3.3.3), the polynomial $A_{\boldsymbol{\mu}}(t)$ must be non-zero and so, by Theorem 3.2.2, the dimension vector $\mathbf{v}_{\boldsymbol{\mu}}$ is a root.

Let us now outline the proof of the positivity which is similar to the proof of the main result of [7]. Denote by $\mathbb{K}$ an arbitrary algebraic closure of $\mathbb{F}_q$ and put $\mathfrak{gl}_n:=\mathfrak{gl}_n(\mathbb{K})$. Denote by $F:\mathfrak{gl}_n\to\mathfrak{gl}_n$ the Frobenius endomorphism that raises coefficients of matrices to their q-th power. Say that a tuple $(C_1,\dots,C_k)$ of adjoint orbits of $\mathfrak{gl}_n$ is *generic* [11, §5.1] if $\sum_{i=1}^k\mathrm{Tr}\,(C_i)=0$ and if for any subspace $V\subset\mathbb{K}^n$ stable by some $X_i\in C_i$ for each $i=1,\dots,k$, such that

$$\sum_{i=1}^k\mathrm{Tr}\,(X_i|_V)=0$$

then either $V=0$ or $V=\mathbb{K}^n$. Generic tuples of semisimple regular adjoint orbits always exists [5, §2.2][11, §5.1]. Recall that the G-conjugacy classes of F-stable maximal tori of $\mathrm{GL}_n=\mathrm{GL}_n(\mathbb{K})$ are parametrized by the conjugacy classes of $\mathfrak{S}_n$. For w in the symmetric group $\mathfrak{S}_n$ we denote by T_w a representative of the corresponding G-conjugacy class of maximal tori. Say that an F-stable regular semisimple adjoint orbit of $\mathfrak{gl}_n$ is of type $w\in\mathfrak{S}_n$ if it has a non-empty intersection with $\mathfrak{t}_w^F$ where $\mathfrak{t}_w:=\mathrm{Lie}(T_w)$. Denote by $\mathbb{S}_n=\mathfrak{S}_n\times\cdots\times\mathfrak{S}_n$ the Cartesian product of k copies of $\mathfrak{S}_n$. Now for each conjugacy class of $\mathbb{S}_n$ with representative $\mathbf{w}=(w_1,\dots,w_k)$ choose a generic tuple $(C^{w_1},\dots,C^{w_k})$ of F-stable semisimple regular adjoint orbits of $\mathfrak{gl}_n$ of type $\mathbf{w}$ (such a choice is possible for any $\mathbf{w}$ assuming that q is sufficiently large which we now assume). Consider the space

$$\mathcal{V}^{\mathbf{w}}:=\left\{(A_1,B_1,\dots,A_g,B_g,X_1,\dots,X_k)\in\mathfrak{gl}_n^{2g}\times C^{w_1}\times\cdots\times C^{w_k}\;\middle|\;\sum_i[A_i,B_i]+\sum_jX_j=0\right\},$$

and the affine GIT quotient

$$\mathcal{Q}^{\mathbf{w}}:=\mathcal{V}^{\mathbf{w}}/\!/\mathrm{GL}_n:=\mathrm{Spec}\left(\mathbb{K}[\mathcal{V}^{\mathbf{w}}]^{\mathrm{GL}_n}\right)$$

where GL_n acts on $\mathcal{V}^{\mathbf{w}}$ diagonally by conjugation. It is well-known (see for instance [5]) that $\mathcal{Q}^{\mathbf{w}}$ is non-singular, irreducible and has vanishing odd cohomology. If $\mathbf{w}=1$, we will simply write $\mathcal{Q}$ instead of $\mathcal{Q}^1$. We know by Lemma 7.2.1, Theorem 6.9.1 and Theorem 6.10.1 in [11] that

$$\epsilon(\mathbf{w}_\lambda)\;\#\mathcal{Q}^{\mathbf{w}_\lambda}(\mathbb{F}_q)=q^{d/2}\,\langle\mathbb{V}(q),p_\lambda\rangle\tag{3.3.4}$$

where

$$d := n^2(2g - 2 + k) - kn + 2$$

is the dimension of Q, ϵ is the sign character of $\mathbb{S}_n$ and where $\mathbf{w}_\lambda \in \mathbb{S}_n$ is in the conjugacy class corresponding to the multi-partition $\boldsymbol{\lambda} \in \overline{\mathcal{P}}$. By definition $X_\lambda(t) := \langle \mathbb{V}(t), p_\lambda \rangle$ is a rational function in t and by (3.3.4) the function $t^{d/2}X_\lambda(t)$ is an integer for infinitely many values of t, therefore $t^{d/2}X_\lambda(t)$ must by a polynomial in t with rational coefficients.

On the other hand, exactly as in [7, §2.2.1] we can prove (assuming that the characteristic is large enough) that there exists, for each i, a representation $\rho^i : \mathbb{S}_n \rightarrow \mathrm{GL}(H_c^{2i}(Q))$, where $H_c^{2i}(Q)$ denotes the compactly supported ℓ-adic cohomology of Q, such that

$$\#Q^{\mathbf{w}}(\mathbb{F}_q) = \sum_{i=d/2}^{d} \mathrm{Trace}\,\left(\rho^i(\mathbf{w})\,|\,H_c^{2i}(Q)\right) q^i. \tag{3.3.5}$$

Note that the character $\psi^i : \mathbb{S}_n \to \mathbb{Q}$, $\mathbf{w} \mapsto \mathrm{Trace}\,\left(\rho^i(\mathbf{w})\,|\,H_c^{2i}(Q)\right)$ of the representation ρ^i does not depend on q. *A priori* it should depend on the characteristic of $\mathbb{K}$ but it does not because it follows from the identities (3.3.4) and (3.3.5) that the values of the characters of the representations ρ^i are given by the coefficients of the polynomials $t^{d/2}X_\lambda(t) \in \mathbb{Q}[t]$. If $Q/\mathbb{C}$ denotes the complex analogue of Q (see 1.1.1), then we know by [15, see below Lemma 48] that there is a representation $\rho^i_{\mathbb{C}}$ of $W_{\mathbf{v}}$ on $H_c^{2i}(Q/\mathbb{C}, \mathbb{C})$. We can actually prove (as in [7, §2.2.1]) that ψ^i is also the character of $\rho^i_{\mathbb{C}}$. From the identity $s_\mu = \sum_\lambda z_\lambda^{-1}\chi^\mu_\lambda p_\lambda$ we find that

$$V_\mu(q) = \sum_\lambda z_\lambda^{-1}\chi^\mu_\lambda \langle \mathbb{V}(q), p_\lambda \rangle .$$

Combining Formulas (3.3.5) and (3.3.4) we get that

$$\begin{aligned} V_\mu(q) &= q^{-d/2} \sum_i \left(\sum_\lambda z_\lambda^{-1}\chi^\mu_\lambda \epsilon(\mathbf{w}_\lambda)\psi^i(\mathbf{w}_\lambda) \right) q^i \\ &= q^{-d/2} \sum_i \left\langle \chi^{\mu'}, \psi^i \right\rangle_{\mathbb{S}_n} q^i, \end{aligned}$$

where $\boldsymbol{\mu}'$ denote the dual multi-partition of $\boldsymbol{\mu}$. Since these identities are true for infinitely many values of q, the coefficients of the polynomial $V_\mu(t)$ coincide with the multiplicities $\left\langle \chi^{\mu'}, \psi^i \right\rangle_{\mathbb{S}_n}$ and therefore are non-negative integers. □

3.4 The unipotent case

For $\boldsymbol{\mu} \in \overline{\mathcal{P}} - \{0\}$, define

$$U_\mu(t) := \left\langle \mathbb{U}(t), s_\mu \right\rangle$$

where $\mathbb{U}(t)$ is as in §2.5 and put $U_\mu(t) := 1$ if $\boldsymbol{\mu} = 0$.

Proposition 3.4.1. *For all $\boldsymbol{\mu} \in \overline{\mathcal{P}} - \{0\}$, we have*

$$U_\mu(q) = \left\langle \mathcal{E} \otimes \mathcal{U}_{\mu^1} \otimes \cdots \otimes \mathcal{U}_{\mu^k}, 1 \right\rangle . \tag{3.4.1}$$

Recall the following relation between $V_{\boldsymbol{\mu}}(t)$ and $U_{\boldsymbol{\mu}}(t)$,

$$\sum_{\boldsymbol{\mu}} U_{\boldsymbol{\mu}}(t) s_{\boldsymbol{\mu}} = \mathrm{Exp}\left(\sum_{\boldsymbol{\mu}} V_{\boldsymbol{\mu}}(t) s_{\boldsymbol{\mu}} \right). \tag{3.4.2}$$

To prove Proposition 3.4.1 we recall the definition of the type of a conjugacy class of G. Let $F : \overline{\mathbb{F}}_q \to \overline{\mathbb{F}}_q$, $x \mapsto x^q$ be the Frobenius endomorphism and let O be the set of $\langle F \rangle$-orbits of $\overline{\mathbb{F}}_q^{\times}$. The conjugacy classes of G corresponds to the maps $f : O \to \mathcal{P}$ such that

$$\sum_{\gamma \in O} |\gamma| \cdot |f(\gamma)| = n.$$

Let C be a conjugacy class of G corresponding to such a function f. The type of C is the function $\omega_C = \omega_f \in \mathbf{T}_n$ defined by

$$\omega_C(d, \lambda) := \# \{ \gamma \in O \,|\, (d, \lambda) = (|\gamma|, f(\gamma)) \}.$$

Proof of Proposition 3.4.1. We have

$$\begin{aligned} \left\langle \mathcal{E} \otimes \mathcal{U}_{\mu^1} \otimes \cdots \otimes \mathcal{U}_{\mu^k}, 1 \right\rangle &= \frac{1}{|G|} \sum_{g \in G} \mathcal{E}(g) \mathcal{U}_{\mu^1}(g) \cdots \mathcal{U}_{\mu^k}(g) \\ &= \sum_C \frac{\mathcal{E}(C)}{a_C(q)} \prod_{i=1}^k \mathcal{U}_{\mu^i}(C) \end{aligned}$$

where the last sum is over the conjugacy classes of G and where $a_C(q)$ denotes the cardinality of the centralizer in G of an element of C. It is well-known (see for instance [6, Theorem 2.2.2]) that for any conjugacy class C of G and any partition μ of n we have

$$\mathcal{U}_\mu(C) = \left\langle \tilde{H}_{\omega_C}(\mathbf{x}; q), s_\mu(\mathbf{x}) \right\rangle.$$

By Formula (2.5.1) we also have

$$\mathcal{H}_{\omega_C}(q) = \frac{\mathcal{E}(C)}{a_C(q)}.$$

Let $\mathbf{C}_n$ be the set of conjugacy classes of $\mathrm{GL}_n(\mathbb{F}_q)$ and put $\mathbf{C} = \cup_{n \geq 1} \mathbf{C}_n$. Denote also by $\mathcal{P}^O$ the set of all function $O \to \mathcal{P}$ with finite support. If 0 denotes the function that take the value 0 everywhere, we put $\mathcal{H}_{\omega_0}(q) = \tilde{H}_{\omega_0}(\mathbf{x}; q) = 1$.

$$\begin{aligned} \sum_{\boldsymbol{\mu} \in \overline{\mathcal{P}}} \left\langle \mathcal{E} \otimes \mathcal{U}_{\mu^1} \otimes \cdots \otimes \mathcal{U}_{\mu^k}, 1 \right\rangle s_{\boldsymbol{\mu}} &= 1 + \sum_{C \in \mathbf{C}} \mathcal{H}_{\omega_C}(q) \prod_i \tilde{H}_{\omega_C}(\mathbf{x}_i; q) \\ &= \sum_{f \in \mathcal{P}^O} \mathcal{H}_{\omega_f}(q) \prod_i \tilde{H}_{\omega_f}(\mathbf{x}_i; q) \\ &= \prod_{\gamma \in O} \Omega\left(\mathbf{x}_1^{|\gamma|}, \ldots \mathbf{x}_k^{|\gamma|}; q^{|\gamma|} \right) \\ &= \prod_{d=1}^{\infty} \Omega\left(\mathbf{x}_1^d, \ldots, \mathbf{x}_k^d; q^d \right)^{\phi_d(q)} \end{aligned}$$

where $\phi_d(q)$ denotes the number of elements in O of size d. Recall that

$$\phi_n(q)=\frac{1}{n}\sum_{d|n}\mu(d)(q^{n/d}-1).$$

By Proposition 2.1.2 we deduce that

$$\mathrm{Log}\left(\sum_{\mu\in\overline{\mathcal{P}}}\left\langle\mathcal{E}\otimes\mathcal{U}_{\mu^1}\otimes\cdots\otimes\mathcal{U}_{\mu^k},1\right\rangle s_{\mu}\right)=(q-1)\,\mathrm{Log}\,\Omega(q).$$

□

From Proposition 3.4.1, Theorem 2.4.4 and Theorem 3.3.2(2) we deduce the following one.

Theorem 3.4.2. *Let* $\boldsymbol{\mu}\in\overline{\mathcal{P}}-\{0\}$. *We have the following assertions.*
(i) The polynomial $U_{\boldsymbol{\mu}}(t)$ *has non-negative integer coefficients.*
(ii) The polynomial $U_{\boldsymbol{\mu}}(t)$ *is non-zero if and only if there exists* $\boldsymbol{\omega}^o=\boldsymbol{\alpha}_1^{n_1}\cdots\boldsymbol{\alpha}_s^{n_s}\in\overline{\mathbf{T}}^o$ *and* $(\tau^1,\dots,\tau^s)\in\mathcal{R}_{\omega^o,\boldsymbol{\mu}}$ *such that for all* $i=1,\dots,s$ *we have*

$$\ell(\tau^i)\leq V_{\alpha_i}(1). \tag{3.4.3}$$

Remark 3.4.3. Note that, by Theorem 3.3.2 (1), the inequality (3.4.3) does not hold unless $\mathbf{v}_{\alpha_i}$ is a root of Γ_{α_i}.

We are now going to give a simple sufficient condition for $U_{\boldsymbol{\mu}}(t)$ to be non-zero.

Denote by $\overline{\mathbf{T}}^o_+$ the set of $\boldsymbol{\omega}^o=\boldsymbol{\alpha}_1^{n_1}\cdots\boldsymbol{\alpha}_s^{n_s}\in\overline{\mathbf{T}}^o$ such that $\mathbf{v}_{\alpha_i}$ is root of Γ_{α_i}. By Proposition 3.4.1, §2.4 and Remark 3.4.3 we have a decomposition

$$U_{\boldsymbol{\mu}}(t)=\sum_{\omega^o\in\overline{\mathbf{T}}^o_+}W_{\boldsymbol{\mu}}^{\omega^o}(t) \tag{3.4.4}$$

where

$$W_{\boldsymbol{\mu}}^{\omega^o}(t)=\sum_{\omega\in\overline{\mathfrak{H}}^{-1}(\omega^o)}A^o_{\omega}V_{\omega}(t)c^{\boldsymbol{\mu}}_{\omega}.$$

For $\boldsymbol{\omega}^o=\boldsymbol{\alpha}_1^{n_1}\cdots\boldsymbol{\alpha}_s^{n_s}\in\overline{\mathbf{T}}^o_+$, put

$$d_{\omega^o}:=\sum_{i=1}^{s}n_id_{\alpha_i}$$

where d_{α}, with $\boldsymbol{\alpha}\in\overline{\mathcal{P}}$, is given by Formula (3.2.3).

For all $\boldsymbol{\omega}\in\overline{\mathfrak{H}}^{-1}(\boldsymbol{\omega}^o)$, the degree of the polynomial $V_{\omega}(t)$ is $d_{\omega^o}/2$ by Theorem 3.3.2 (2).

By Remark 2.4.2 we have

$$\sum_{\omega\in\overline{\mathfrak{H}}^{-1}(\omega^o)}A^o_{\omega}c^{\boldsymbol{\mu}}_{\omega}=\left\langle\mathbb{C}^{\boldsymbol{\mu}}_{\omega^o},1\right\rangle_{W_{\omega^o}}.$$

We deduce the following proposition.

Proposition 3.4.4. *Let* $\omega^o=\alpha_1^{n_1}\cdots\alpha_s^{n_s}\in\overline{\mathbf{T}}^o_+$. *If* $\left\langle\mathbb{C}^{\boldsymbol{\mu}}_{\omega^o},1\right\rangle_{W_{\omega^o}}\neq 0$, *then*

$$\left\langle\mathbb{C}^{\boldsymbol{\mu}}_{\omega^o},1\right\rangle_{W_{\omega^o}}q^{d_{\omega^o}}$$

is the term of $W^{\omega^o}_{\boldsymbol{\mu}}(t)$ *of highest degree. In particular, if for all* $i = 1, \dots, s$ *the vector* $\mathbf{v}_{\alpha_i}$ *is a real root (in which case* $d_{\alpha_i} = 0$*), then*

$$W^{\omega^o}_{\boldsymbol{\mu}}(t) = \left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle_{W_{\omega^o}}.$$

Remark 3.4.5. Let $\boldsymbol{\mu} = (\mu^1, \dots, \mu^k) \in \overline{\mathcal{P}}$ and $\boldsymbol{\omega}^o = \boldsymbol{\alpha}_1^{n_1} \cdots \boldsymbol{\alpha}_s^{n_s} \in \overline{\mathbf{T}}^o$ be of same size n. Notice that if $\boldsymbol{\alpha}_i = ((1), \dots, (1))$ for all i, then $W_{\omega^o} = \mathfrak{S}_n$ and

$$\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle = \langle H_{\boldsymbol{\mu}}, 1 \rangle$$

where $H_{\boldsymbol{\mu}}$ is the $\mathbb{C}[\mathfrak{S}_n]$-module $H_{\mu^1} \otimes \cdots \otimes H_{\mu^k}$. Note also that if $n_1 = n_2 = \cdots = n_s = 1$ then $W_{\omega^o} = 1$ and

$$\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle = \dim \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}.$$

Hence in this case $\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle$ is a product of Littlewood-Richardson coefficients. If moreover $s = 1$ and $n_1 = 1$, then

$$\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle = \delta_{\omega^o, \boldsymbol{\mu}}.$$

Corollary 3.4.6. *Let* $\boldsymbol{\mu} \in \overline{\mathcal{P}} - \{0\}$*. If there exists* $\boldsymbol{\omega}^o \in \overline{\mathbf{T}}^o_+$ *such that* $\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle \neq 0$*, then* $U_{\boldsymbol{\mu}}(t) \neq 0$*. In particular, if* $\mathbf{v}_{\boldsymbol{\mu}}$ *is a root or if* $\left\langle H_{\boldsymbol{\mu}}, 1 \right\rangle \neq 0$*, then* $U_{\boldsymbol{\mu}}(t) \neq 0$*.*

Proof. By Theorem 3.4.2 (i), the polynomials $W^{\omega^o}_{\boldsymbol{\mu}}(t)$ have non-negative integer coefficients and so there are no cancellation in the decomposition (3.4.4). We can now apply Proposition 3.4.4 to deduce that if $\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle \neq 0$ for some $\boldsymbol{\omega}^o$, then $U_{\boldsymbol{\mu}}(t) \neq 0$. □

Note that Corollary 3.4.6 is also a straightforward consequence of Theorem 3.4.2. Indeed if $\left\langle \mathbb{C}^{\boldsymbol{\mu}}_{\omega^o}, 1 \right\rangle \neq 0$ with $\boldsymbol{\omega}^o = \boldsymbol{\alpha}_1^{n_1} \cdots \boldsymbol{\alpha}_s^{n_s} \in \overline{\mathbf{T}}^o_+$, then the multi-partition $((n_1)^1, \dots, (n_s)^1)$ belongs to $\mathcal{R}_{\omega^o, \boldsymbol{\mu}}$ and clearly $\ell((n_i)^1) = 1 \leq V_{\alpha_i}(1)$ as $\boldsymbol{\alpha}_i$ is a root.

Theorem 3.4.7. *Let* $\boldsymbol{\mu} \in \overline{\mathcal{P}} - \{0\}$*.*

(i) If $\mathbf{v}_{\boldsymbol{\mu}}$ *is a root of* $\Gamma_{\boldsymbol{\mu}}$*, then the degree of the polynomial* $U_{\boldsymbol{\mu}}(t)$ *is at least* $d_{\boldsymbol{\mu}}/2$*.*

(ii) If $\delta(\boldsymbol{\mu}) \geq 2$*, then the degree of* $U_{\boldsymbol{\mu}}(t)$ *is exactly* $d_{\boldsymbol{\mu}}/2$*.*

(ii) If $\delta(\boldsymbol{\mu}) \geq 3$ *or* $g = 0$*,* $k = 3$ *and* $\delta(\boldsymbol{\mu}) = 2$*, then* $U_{\boldsymbol{\mu}}(t)$ *is a monic polynomial.*

Remark 3.4.8. The degree of $U_{\boldsymbol{\mu}}(t)$ may be strictly larger than $d_{\boldsymbol{\mu}}/2$. Indeed, assume that $g = 0$, $n = 6$ and $k = 3$, and take $\boldsymbol{\mu} = ((2^3), (2^3), (2^3))$. Note that $\mathbf{v}_{\boldsymbol{\mu}} = 2 \cdot \boldsymbol{\alpha}_1$ where $\boldsymbol{\alpha}_1$ is the indivisible imaginary root of $\tilde{E}_6$. A direct calculation shows that $d_{\boldsymbol{\mu}} = 2 = d_{\alpha_1}$. Consider $\boldsymbol{\omega}^o = \boldsymbol{\alpha}_1^2$. Then a direct calculation, using that $V_{\alpha_1}(t) = t$ (see Section §3.5) shows that $W^{\omega^o}_{\boldsymbol{\mu}}(t) = t^2$ while $d_{\boldsymbol{\mu}}/2 = 1$.

For $\boldsymbol{\mu} = (\mu^1, \dots, \mu^k) \in \overline{\mathcal{P}}$, put

$$\Delta(\boldsymbol{\mu}) := \frac{1}{2} d_{\boldsymbol{\mu}} - 1 = \frac{1}{2}(2g - 2 + k)n^2 - \frac{1}{2} \sum_{i,j} (\mu^i_j)^2.$$

Notice that if $\boldsymbol{\alpha} = (\alpha^1, \dots, \alpha^k) \in \overline{\mathcal{P}}$ (possibly of size different from $|\boldsymbol{\mu}|$), then

$$2|\boldsymbol{\mu}|\Delta(\boldsymbol{\alpha}) = \delta(\boldsymbol{\mu})|\boldsymbol{\alpha}|^2 + \sigma_{\boldsymbol{\mu}}(\boldsymbol{\alpha}).$$

where $\sigma_{\boldsymbol{\mu}}(\boldsymbol{\alpha}) := \sum_{i=1}^k \sigma_{\mu^i}(\alpha^i)$.

Proposition 3.4.9. *Let* $\boldsymbol{\mu}, \boldsymbol{\alpha}_1, \ldots, \boldsymbol{\alpha}_s \in \overline{\mathcal{P}} - \{0\}$ *such that* $|\boldsymbol{\mu}| = |\sum_{i=1}^s \boldsymbol{\alpha}_i|$ *and* $\boldsymbol{\mu} \trianglelefteq \sum_{i=1}^s \boldsymbol{\alpha}_i$.

(i) Assume that for all $i = 1, \ldots, s$, *we have* $\delta(\boldsymbol{\mu})|\boldsymbol{\alpha}_i| \geq 2$. *Then*

$$d_{\boldsymbol{\mu}} \geq \sum_i d_{\boldsymbol{\alpha}_i} \tag{3.4.5}$$

If moreoever $\delta(\boldsymbol{\mu})|\boldsymbol{\alpha}_i| > 2$ *for some* $i = 1, \ldots, s$, *then the inequality (3.4.5) is strict.*

(ii) Assume that $g = 0$, $k = 3$ *and* $\delta(\boldsymbol{\mu}) \geq 2$. *Then the inequality (3.4.5) is strict.*

Proof. Put $n := |\boldsymbol{\mu}|$. Let us first prove (i).

$$2n\Delta(\boldsymbol{\mu}) - 2n\sum_{i=1}^s \Delta(\boldsymbol{\alpha}_i) = \delta(\boldsymbol{\mu})\left(n^2 - \sum_i |\boldsymbol{\alpha}_i|^2\right) + \sigma_{\boldsymbol{\mu}}(\boldsymbol{\mu}) - \sum_{i=1}^s \sigma_{\boldsymbol{\mu}}(\boldsymbol{\alpha}_i).$$

We thus have

$$\begin{aligned} nd_{\boldsymbol{\mu}} - n\sum_{i=1}^s d_{\boldsymbol{\alpha}_i} &= 2n\Delta(\boldsymbol{\mu}) - 2n\sum_{i=1}^s \Delta(\boldsymbol{\alpha}_i) - 2n(s-1) \\ &= \delta(\boldsymbol{\mu})\left(\sum_{i \neq j} |\boldsymbol{\alpha}_i|\,|\boldsymbol{\alpha}_j|\right) - 2n(s-1) + \sigma_{\boldsymbol{\mu}}(\boldsymbol{\mu}) - \sum_{i=1}^s \sigma_{\boldsymbol{\mu}}(\boldsymbol{\alpha}_i) \\ &= \sum_{i=1}^s \sum_{j \neq i} |\boldsymbol{\alpha}_i| \left(\delta(\boldsymbol{\mu})|\boldsymbol{\alpha}_j| - 2\right) + \sigma_{\boldsymbol{\mu}}(\boldsymbol{\mu}) - \sum_{i=1}^s \sigma_{\boldsymbol{\mu}}(\boldsymbol{\alpha}_i) \end{aligned}$$

By Theorem 2.6.1 we have $\sigma_{\boldsymbol{\mu}}(\boldsymbol{\mu}) - \sum_{i=1}^s \sigma_{\boldsymbol{\mu}}(\boldsymbol{\alpha}_i) \geq 0$ hence the assertion (i).

We now prove (ii). If $|\boldsymbol{\alpha}| \leq 2$ then a straightforward calculation shows that $d_{\boldsymbol{\alpha}} \leq 0$, hence if for all $i = 1, \ldots, s$, we have $|\boldsymbol{\alpha}_i| \leq 2$, then clearly $d_{\boldsymbol{\mu}} - \sum_i d_{\boldsymbol{\alpha}_i}$ is strictly positive.

□

Proof of Theorem 3.4.7. By the decomposition (3.4.4), we have

$$U_{\boldsymbol{\mu}}(t) = V_{\boldsymbol{\mu}}(t) + \sum_{\omega^o \neq \boldsymbol{\mu}} W_{\boldsymbol{\mu}}^{\omega^o}(t).$$

Recall that when $\mathbf{v}_{\boldsymbol{\mu}}$ is a root, the polynomial $V_{\boldsymbol{\mu}}(t)$ is non-zero monic of degree $d_{\boldsymbol{\mu}}/2$ by Theorem 3.3.2. Hence (i). The assumption $\delta(\boldsymbol{\mu}) \geq 2$ implies (by Proposition 3.4.9) that the degree of $W_{\boldsymbol{\mu}}^{\omega^o}(t)$ is smaller or equal to $d_{\boldsymbol{\mu}}/2$. Since the leading coefficients of $W_{\boldsymbol{\mu}}^{\omega^o}(t)$ are non-negative we get assertion (ii). The assertion (iii) is also a consequence of Proposition 3.4.9. Indeed, in this case the degrees of $W_{\boldsymbol{\mu}}^{\omega^o}(t)$ are strictly smaller than $d_{\boldsymbol{\mu}}/2$.

□

Corollary 3.4.10. *Denote by* St_n *the Steinberg character of* $\mathrm{GL}_n(\mathbb{F}_q)$. *Then* $\langle \mathrm{St}_n \otimes \mathrm{St}_n \otimes \mathrm{St}_n, 1\rangle$ *is a monic polynomial of degree* $\frac{1}{2}(n-1)(n-2)$ *for all* $n \geq 1$.

Proof. We have $\langle \mathrm{St}_n \otimes \mathrm{St}_n \otimes \mathrm{St}_n, 1\rangle = U_{\boldsymbol{\mu}}(q)$ with $\boldsymbol{\mu} = ((1^n), (1^n), (1^n))$. We find that $d_{\boldsymbol{\mu}} = (n-1)(n-2)$. Now $\delta(\boldsymbol{\mu}) = n - 3$. Hence by Theorem 3.4.7 the corollary is true for $n \geq 5$. The cases $n = 2, 3, 4$ are not difficult to work out with the above results (see §3.5 to see how to apply the above results) and we find that

$$\begin{aligned} \langle \mathrm{St}_2 \otimes \mathrm{St}_2 \otimes \mathrm{St}_2, 1\rangle &= 1 \\ \langle \mathrm{St}_3 \otimes \mathrm{St}_3 \otimes \mathrm{St}_3, 1\rangle &= q + 1 \\ \langle \mathrm{St}_4 \otimes \mathrm{St}_4 \otimes \mathrm{St}_4, 1\rangle &= q^3 + 2q + 1 \end{aligned}$$

Note that for $n \geq 3$, we have $\delta(\boldsymbol{\mu}) \geq 0$ and so $\mathbf{v}_\mu$ is in the fundamental domain of imaginary root (see Proposition 3.2.1). For $n = 2$, the vector $\mathbf{v}_\mu$ is a real root of D_4.

□

3.5 Example

Assume that $n = k = 3$ and $g = 0$. We first need to list the $V_\alpha(t)$ where $|\alpha| \leq 3$ and $\mathbf{v}_\alpha \in \Phi(\Gamma_\alpha)$. This is given in the left column of the table below (up to permutation of the coordinates of $\boldsymbol{\alpha}$).

$\boldsymbol{\alpha}$	$\lvert\boldsymbol{\alpha}\rvert$	Γ_α	$\mathbf{v}_\alpha$	$V_\alpha(t)$
$\boldsymbol{\alpha}_1 := ((1^3),(1^3),(1^3))$	3	$\tilde{E}_6$	$(3,(2,1),(2,1),(2,1))$	t
$\boldsymbol{\alpha}_{2,1} := ((2,1),(1^3),(1^3))$	3	E_6	$(3,(1),(2,1),(2,1))$	1
$\boldsymbol{\alpha}_3 := ((1^2),(1^2),(1^2))$	2	D_4	$(2,(1),(1),(1))$	1
$\boldsymbol{\alpha}_4 := ((1),(1),(1))$	1	A_1	(1)	1

(3.5.1)

where in the second column we only put the underlying graph of Γ_α (as the orientation does not matter) and where $\mathbf{v}_\alpha$ is written in the form $(v_0,(v_{[1,1]},v_{[1,2]},\dots),(v_{[2,1]},v_{[2,2]},\dots),\dots,(v_{[k,1]},v_{[k,2]},\dots))$. Note that $\mathbf{v}_{\alpha_{2,1}}$, $\mathbf{v}_{\alpha_3}$ and $\mathbf{v}_{\alpha_4}$ of the last three rows are real roots and so by Theorem 3.3.2 we have $V_\alpha(t) = 1$ in these cases. Notice that $\mathbf{v}_{\alpha_1}$ of the first row is the unique indivisible positive imaginary root of $\tilde{E}_6$. We can see that $V_{\alpha_1}(t) = t$ either by computing directly $V_{\alpha_1}(t)$ with Formula (3.4.1) or by proceeding as follows. Put $\boldsymbol{\alpha}_{2,2} := ((1^3),(2,1),(1^3))$ and $\boldsymbol{\alpha}_{2,3} := ((1^3),(1^3),(2,1))$. Applying Formula (3.3.3) we find that

$$\begin{aligned} A_{\alpha_1}(t) &= V_{\alpha_1}(t) + \sum_{i=1}^{3} 2\, V_{\alpha_{2,i}}(t) \\ &= V_{\alpha_1}(t) + 6 \end{aligned}$$

On the other hand the polynomial $A_{\alpha_1}(t)$ is monic of degree 1 by Theorem 3.2.2 and we know by [1] that the constant term of the polynomials $A_{\alpha_1}(t)$ is the multiplicity of the root $\mathbf{v}_{\alpha_1}$. Now the multiplicity of $\mathbf{v}_{\alpha_1}$ is 6 (see [10, Chap. 7, Corollary 7.4]) and so $A_{\alpha_1}(t) = t + 6$. Hence $V_{\alpha_1}(t) = t$.

The set of types $\boldsymbol{\omega}^o \in \overline{\mathbf{T}}^o_+$ of size 3 are thus

$$\boldsymbol{\alpha}_1,\ \boldsymbol{\alpha}_{2,1},\ \boldsymbol{\alpha}_{2,2},\ \boldsymbol{\alpha}_{2,3},\ \boldsymbol{\alpha}_3\boldsymbol{\alpha}_4,\ \boldsymbol{\alpha}_4^3.$$

and so the decomposition (3.4.4) reads

$$U_\mu(t) = W_\mu^{\alpha_1}(t) + W_\mu^{\alpha_{2,1}}(t) + W_\mu^{\alpha_{2,2}} + W_\mu^{\alpha_{2,3}} + W_\mu^{\alpha_3\alpha_4}(t) + W_\mu^{\alpha_4^3}(t).$$

Since the dimension vectors associated to $\boldsymbol{\alpha}_{2,1}$, $\boldsymbol{\alpha}_{2,2}$, $\boldsymbol{\alpha}_{2,3}$, $\boldsymbol{\alpha}_3$, $\boldsymbol{\alpha}_4$, $\boldsymbol{\alpha}_4$ are real roots, by Proposition 3.4.4 and Remark 3.4.5 we have

$$\begin{aligned} U_\mu(t) &= \delta_{\alpha_1,\mu} V_{\alpha_1}(t) + \delta_{\alpha_{2,1},\mu} + \delta_{\alpha_{2,2},\mu} + \delta_{\alpha_{2,3},\mu} + c^{\boldsymbol{\mu}}_{\mathbf{(1^2)(1)}} + \left\langle H_\mu, 1 \right\rangle \\ &= \delta_{\alpha_1,\mu} t + \delta_{\alpha_{2,1},\mu} + \delta_{\alpha_{2,2},\mu} + \delta_{\alpha_{2,3},\mu} + c^{\boldsymbol{\mu}}_{\mathbf{(1^2)(1)}} + \left\langle H_\mu, 1 \right\rangle \end{aligned}$$

where $c^{\boldsymbol{\mu}}_{\mathbf{(1^2)(1)}} := c^{\mu^1}_{(1^2)(1)} c^{\mu^2}_{(1^2)(1)} c^{\mu^3}_{(1^2)(1)}$ with $(\mu^1,\mu^2,\mu^3) = \boldsymbol{\mu}$.

Recall that $c^{\mu}_{(1^2)(1)} = 1$ if $\mu \in \{(2,1),(1,1,1)\}$ and $c^{\mu}_{(1^2)(1)} = 0$ otherwise.

Hence

$\boldsymbol{\mu}$	$\delta_{\boldsymbol{\alpha}_1,\boldsymbol{\mu}}$	$\delta_{\boldsymbol{\alpha}_{2,1},\boldsymbol{\mu}}$	$c^{\boldsymbol{\mu}}_{(\mathbf{1}^2)(\mathbf{1})}$	$\left\langle H_{\boldsymbol{\mu}}, 1\right\rangle$	$U_{\boldsymbol{\mu}}(t)$
$((3^1),(3^1),(3^1))$	0	0	0	1	1
$((3^1),(3^1),(2,1))$	0	0	0	0	0
$((3^1),(3^1),(1^3))$	0	0	0	0	0
$((3^1),(2,1),(2,1))$	0	0	0	1	1
$((3^1),(2,1),(1^3))$	0	0	0	0	0
$((3^1),(1^3),(1^3))$	0	0	0	1	1
$((2,1),(2,1),(2,1))$	0	0	1	1	2
$((2,1),(2,1),(1^3))$	0	0	1	1	2
$((2,1),(1^3),(1^3))$	0	1	1	0	2
$((1^3),(1^3),(1^3))$	1	0	1	0	$t+1$

References

[1] Crawley-Boevey, W. and Van den Bergh, M.: Absolutely indecomposable representations and Kac-Moody Lie algebras. With an appendix by Hiraku Nakajima. *Invent. Math.* **155** (2004), no. 3, 537–559.

[2] Garsia, A.M. and Haiman, M.: A remarkable q,t-Catalan sequence and q-Lagrange inversion, *J. Algebraic Combin.* **5** (1996) no. 3, 191-244.

[3] Getzler, E.: Mixed Hodge Structures of configuration spaces, preprint, arXiv:alg-geom/9510018.

[4] Green, J.A.: The characters of the finite general linear groups. *Trans. Amer. Math. Soc.* **80** (1955), 402–447.

[5] Hausel T., Letellier, E. and Rodriguez-Villegas, F.: Arithmetic harmonic analysis on character and quiver varieties, *Duke Math. J.*, **160** (2011) no. 2, 323–400.

[6] Hausel T., Letellier, E. and Rodriguez-Villegas, F.: Arithmetic harmonic analysis on character and quiver varieties II, preprint arXiv:1109.5202.

[7] Hausel T., Letellier, E. and Rodriguez-Villegas, F.: Positivity of A-polynomials and DT-invariants of quivers, preprint (to be posted on arXiv soon).

[8] Hiss, Gerhard and Lübeck, Frank: Some observations on products of characters of finite classical groups. *Finite groups 2003*, 195–207, Walter de Gruyter GmbH & Co. KG, Berlin (2004).

[9] Kac, V.: Root systems, representations of quivers and invariant theory. *Lecture Notes in Mathematics*, **vol. 996**, Springer-Verlag (1982), 74–108.

[10] Kac, V.: Infinite dimensional Lie algebras. Third edition. *Cambridge University Press*, Cambridge (1990), xxii+400 pp.

[11] Letellier, E.: Quiver varieties and the character ring of general linear groups over finite fields. To appear in *Journal of the European Math. Soc.*, preprint arXiv:1103.2759.

[12] Lübeck, F.: Generic decomposition of tensor products.

http://www.math.rwth-aachen.de/~Frank.Luebeck/preprints/tprdexmpl/tendec.html

[13] Lusztig, G.: Quiver varieties and Weyl group actions, *Ann. Inst. Fourier (Grenoble)* **50** (2000), 461–489.

[14] Macdonald, I.G: Symmetric Functions and Hall Polynomials, *Oxford Mathematical Monographs, second ed., Oxford Science Publications*. The Clarendon Press Oxford University Press, New York, 1995.

[15] Maffei, A.: A remark on quiver varieties and Weyl group, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **1** (2002), no. 3, 649–686.

[16] Mozgovoy, S.: A computational criterion for the Kac conjecture, *J. Algebra* **318** (2007), no. 2, 669–679.

[17] Nakajima, H.: Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras, *Duke Math. J.* **76** (1994), 365–416.

[18] Nakajima, H.: Reflection functors for quiver varieties and Weyl group actions. *Math. Ann.* **327** (2003), no. 4, 671–721.

[19] Saito, M.: Mixed Hodge Modules. *Publ. Res. Inst. Math. Sci.*, **26** (1990), 221–333.